

07/28/00



3c872 U.S. PRO

UTILITY PATENT APPLICATION TRANSMITTAL (Large Entity)
Only for new nonprovisional applications under 37 C.F.R. 1.53(b)

Docket No.:

2804-J

Express Mail Label No.:

EL591095389US

TO THE ASSISTANT COMMISSIONER FOR PATENTS
BOX PATENT APPLICATION
Washington, D.C. 20231

Transmitted herewith for filing under 35 U.S.C. 111(a) and 37 C.F.R. 1.53(b) is a new utility patent application for an invention entitled:

CD30 LIGAND (as amended)

and invented by:

Raymond Goodwin, Craig Smith, Richard Armitage and Hans-Juergen Gruss.

If a **CONTINUING APPLICATION**, check appropriate box and supply the requisite information:

☐ Continuation ☒ Divisional ☐ Continuation-in-part (CIP)

of prior application No.: 09/079,785

Enclosed are:

Application Elements

1. ☒ Filing fee as calculated and transmitted as described below
2. ☒ Specification including claims and abstract (75 pages total)
3. ☐ Drawing(s); Number of Sheets _____
4. ☒ Oath or Declaration
 - a. ☐ Newly executed
 - b. ☒ Copy from a prior application (37.C.F.R. 1.63(d)) (for continuation/divisional application only)
 - c. ☒ With Power of Attorney ☐ Without Power of Attorney
 - d. ☐ DELETION OF INVENTOR(S)
 Signed statement attached deleting inventor(s) named in prior application, see 37 C.F.R. 1.63(d)(2) and 1.33(b).
5. ☒ Incorporation by Reference (usable if Box 4b is checked)
 The entire disclosure of the prior application from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
6. ☐ Computer Program in Microfiche (Appendix)
7. ☒ Nucleotide and/or Amino Acid Sequence Submission
 - a. ☒ Paper copy
 - ☒ Pages 48 - 70 of specification
 - ☐ Separately numbered pages _____ - _____
 - b. ☐ Computer Readable Copy
 - c. ☐ Statement Verifying Identical Paper and Computer Readable Copy
 - d. ☒ Statement under 37 C.F.R. 1.821(e) in lieu of Computer Readable Copy

Accompanying Application Parts

8. ☐ Assignment
- a. ☐ Executed original Assignment and Recordation Form enclosed
- b. ☒ Prior application is assigned of record to Immunex Corporation
(reel 7066 frame 0094)
9. ☐ 37 C.F.R. 3.73(B) Statement (*when there is an assignee*)
10. ☒ Preliminary Amendment
11. ☒ Acknowledgment postcard
12. ☒ Certificate of Mailing by Express Mail (Label No.: EL591095389US)
13. ☐ Certified Copy of Priority Document(s) (*if foreign priority is claimed*)
14. ☐ Additional Enclosures (*please identify below*):


Fee Calculation and Transmittal

CLAIMS AS FILED (after any Preliminary Amendment submitted herewith)					
For	# Filed	# Allowed	# Extra	Rate	Fee
Total Claims	23	- 20 =	3	x \$18.00	\$54.00
Indep. Claims	4	- 3 =	1	x \$78.00	\$78.00
Multiple Dependent Claims (check if applicable) <input type="checkbox"/>					\$0.00
BASIC FEE					\$690.00
OTHER FEE (<i>specify purpose</i>)					\$0.00
TOTAL FILING FEE					\$822.00

- ☒ The Commissioner is hereby authorized to charge and credit Deposit Account No. 09-0089 as described below. A copy of this sheet is enclosed.
- ☒ Charge the amount of \$822.00 as a filing fee.
- ☒ Credit any overpayment.
- ☒ Charge any additional fees required under 37 C.F.R. 1.16 and 1.17.

Correspondence address:

Immunex Corporation
Law Department
51 University Street
Seattle, Washington 98101
Telephone: (206) 587-0430



Kathryn A. Anderson
Registration No. 32,172

Dated: July 28, 2000

[illegible]

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In the Application of: GOODWIN et al.

Docket No.: 2804-I

Serial No.: -- not yet assigned --

for prior application:

Examiner: L. Spector

Group Art Unit: 1646

Filed: July 28, 2000

For: CD30 LIGAND (as amended)

PRELIMINARY AMENDMENT

Box Patent Application
Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Please amend the above-identified application as follows, prior to examination:

In the Specification:

Page 1, line 7, delete the current title and replace with -- CD30 LIGAND --.

Page 1, line 10, after "This application is" insert -- a divisional of U.S. application Serial No. 09/079,785, filed May 15, 1998, currently allowed, which is a divisional of U.S. application Serial No. 08/580,014, filed December 20, 1995, now U.S. Patent 5,753,203, which is a divisional of U.S. application Serial No. 08/225,989, filed April 12, 1994, now U.S. Patent 5,480,981, which is --.

Page 1, line 11, delete "currently pending" and replace with -- now abandoned -- .

In the Claims:

Cancel claims 1-26. Amend claims 27, 28 and 31 as follows:

27. (amended) A method of delivering a diagnostic or therapeutic agent to CD30⁺ cells, comprising contacting said cells with a conjugate [according to claim 25] comprising a diagnostic or therapeutic agent attached to a CD30 ligand (CD30-L) polypeptide.

28. (amended) A method of delivering a diagnostic or therapeutic agent to CD30⁺ cells, comprising contacting said cells with a conjugate [according to claim 26] comprising a diagnostic or therapeutic agent attached to a CD30-L polypeptide, wherein said CD30-L polypeptide is a soluble fragment of the human CD30-L of SEQ ID NO:8.

31. (amended) A method of treating large cell anaplastic lymphoma (LCAL), comprising administering an effective amount of a soluble CD30-L [according to claim 19] to a human afflicted with LCAL.

Please add new claims 32 to 49, as follows:

32. A method according to claim 27, wherein said CD30-L polypeptide is in the form of an oligomer comprising two or more CD30-L polypeptides, wherein the CD30-L polypeptides are each selected from the group consisting of:

- a) the murine CD30-L of SEQ ID NO:6;
- b) the murine CD30-L of SEQ ID NO:19;
- c) the human CD30-L of SEQ ID NO:8;
- d) the human CD30-L of SEQ ID NO:23; and
- e) a fragment of the CD30-L of (a), (b), (c), or (d), wherein said fragment binds

CD30.

33. A method according to claim 32, wherein said oligomer comprises three CD30-L polypeptides.

34. A method according to claim 32, wherein each of the CD30-L polypeptides is a soluble fragment of the human CD30-L of SEQ ID NO:8 or SEQ ID NO:23.

35. A method according to claim 34, wherein each of the CD30-L polypeptides is selected from the group consisting of a polypeptide comprising the extracellular domain of the human CD30-L of SEQ ID NO:23, and a fragment of said extracellular domain, wherein said fragment binds CD30.

36. A method according to claim 34, wherein said oligomer comprises three CD30-L polypeptides.

37. A method according to claim 35, wherein said oligomer comprises three CD30-L polypeptides.

38. A method according to claim 27, wherein said cells are CD30⁺ lymphoma cells.

39. A method according to claim 28, wherein said cells are CD30⁺ lymphoma cells.

40. A method according to claim 32, wherein said cells are CD30⁺ lymphoma cells.

41. A method according to claim 34, wherein said cells are CD30⁺ lymphoma cells.

42. A method according to claim 36, wherein said cells are CD30⁺ lymphoma cells.
43. A method according to claim 28, wherein said cells are CD30⁺ Hodgkin's Disease cells.
44. A method according to claim 34, wherein said cells are CD30⁺ Hodgkin's Disease cells.
45. A method according to claim 36, wherein said cells are CD30⁺ Hodgkin's Disease cells.
46. A method according to claim 34, wherein said conjugate comprises a therapeutic agent.
47. A method according to claim 31, wherein said soluble CD30-L is a soluble fragment of the human CD30-L of SEQ ID NO:8, wherein said fragment binds CD30.
48. A method for reducing proliferation of LCAL cells in a human afflicted with LCAL, comprising administering an effective amount of a CD30-L oligomer to said human, wherein said oligomer comprises two or three CD30-L polypeptides, wherein each of said CD30-L polypeptides is a soluble fragment of the human CD30-L of SEQ ID NO:8, wherein said oligomer binds CD30.
49. A method according to claim 48, wherein said oligomer comprises three CD30-L polypeptides.

REMARKS

This amendment is submitted in order to place the subject application in better condition for examination. The amendment also serves to add claims directed to particular embodiments of the invention. Each amended and new claim is fully supported by the application as filed.


The title as amended corresponds to the designation employed in the specification, for the subject cytokine. The remaining amendments to page one insert and update information on related applications.

Claims 27, 28, and 31 are amended due to cancellation of the claims on which original claims 27, 28, and 31 were dependent. New claims 32-49 are supported on page 16, line 20, to page 17, line 15; page 4, lines 15-18; page 7, lines 10-20 and 31-33; page 35, line 34, through page 36, line 1; page 2, lines 35-36; page 3, lines 28-32; and page 14, line 16, to

page 16, line 5 of the present specification as originally filed. Further support is found in the language of claims 29 to 31 as originally filed.

Claims 27-49 are now pending in the subject application.

Respectfully submitted,



Kathryn A. Anderson
Registration No. 32,172

Immunex Corporation
51 University Street
Seattle, Washington 98101
(206) 587-0430

IMMUNEX CORPORATION

2804-E

5

TITLE

NOVEL CYTOKINE THAT BINDS CD30

CROSS-REFERENCE TO RELATED APPLICATIONS

- 10 This application is a continuation-in-part of U.S. Application Serial No. 07/966,775, filed on October 27, 1992, currently pending, which is a continuation-in-part of U.S. Application Serial No. 07/907,224, filed on July 1, 1992, now abandoned, which is a continuation-in-part of U.S. Application Serial No. 07/899,660, filed on June 15, 1992, now abandoned, which is a continuation-in-part of U.S.
- 15 Application Serial No. 07/892,459, filed on June 2, 1992, now abandoned, which is a continuation-in-part of U.S. Application Serial No. 07/889,717, filed on May 26, 1992, now abandoned. Priority is also claimed from International Application PCT/US93/04926.

20

BACKGROUND OF THE INVENTION

- Hodgkin's Disease is a human lymphoma, the etiology of which is still not well understood. The neoplastic cells of Hodgkin's Disease are known as Hodgkin and Reed-Sternberg (H-RS) cells. CD30 is a 120 kd surface antigen widely used as a clinical marker for Hodgkin's lymphoma and related hematologic malignancies (Froese et al., *J. Immunol.* 139:2081 (1987); Pfreundschuh et al., *Onkologie* 12:30 (1989); Carde et al., *Eur. J. Cancer* 26:474 (1990)). Originally identified by the monoclonal antibody Ki-1, which is reactive with H-RS cells (Schwab et al., *Nature* (London) 299:65 (1982)), CD30 was subsequently shown to be expressed on a subset of non-Hodgkin's lymphomas (NHL), including Burkitt's lymphoma, as well as several
- 30 virally-transformed lines (human T-Cell Lymphotropic Virus I or II transformed T-cells, and Epstein-Barr Virus transformed B-cells (Stein et al., *Blood* 66:848 (1985); Andreessen et al., *Blood* 63:1299 (1984)). That CD30 plays a role in normal lymphoid interactions is suggested by its histological detection on a small population of lymphoid cells in reactive lymph nodes, and by induced expression on purified T- and B-cells
- 35 following lectin activation (Stein et al., *Int. J. Cancer* 30:445 (1982) and Stein et al., 1985, *supra*).

CD30 expression has also been detected on various non-Hodgkin's lymphomas (NHL), such as large-cell anaplastic lymphomas (LCL), cutaneous T-cell lymphomas, nodular small cleaved-cell lymphomas, lymphocytic lymphomas, peripheral T-cell lymphomas, Lennert's lymphomas, immunoblastic lymphomas, T-cell leukemia/lymphomas (ATLL), adult T-cell leukemia (T-ALL), and centroblastic/centrocytic (cb/cc) follicular lymphomas (Stein et al., *Blood* 66:848 (1985); Miettinen, *Arch. Pathol. Lab. Med.* 116:1197 (1992); Piris et al., *Histopathology* 17:211 (1990); Burns et al., *Am. J. Clin. Pathol.* 93:327 (1990); Piris et al., *Histopathology* 18:25 (1991); Eckert et al., *Am. J. Dermatopathol.* 11:345 (1989); Gianotti et al., *Am. J. Dermatopathol.* 13:503 (1991); Maeda et al., *Br. J. Dermatol.* 121:603 (1989)). The association of the CD30 antigen with lymphoid malignancies has proven to be a useful marker for the identification of malignant cells within lymphoid tissues, particularly lymph nodes. However, expression of CD30 has also been reported on a portion of embryonal carcinomas, nonembryonal carcinomas, malignant melanomas, mesenchymal tumors, and myeloid cell lines and macrophages at late stages of differentiation (Schwartz et al., *Blood* 74:1678 (1989); Pallesen et al., *Am. J. Pathol.* 133:446 (1988); Mechttersheimer et al., *Cancer* 66:1732 (1990); Andreessen et al., *Am. J. Pathol.* 134:187 (1989)).

Cloning and expression of a gene encoding CD30 has been reported and CD30 has been characterized as a transmembrane protein that possesses substantial homology to the nerve growth factor receptor superfamily (Durkop et al., *Cell* 68:421, 1992). Durkop et al. suggest that CD30 is the receptor for one or more as yet unidentified growth factors, and recognize the importance of investigating the existence and nature of such growth factors in order to achieve insight into the etiology of Hodgkin's Disease.

Prior to the present invention, however, no such growth factors or other molecules that bind to the CD30 receptor were known. A need thus remained for identification and characterization of a ligand for CD30.

SUMMARY OF THE INVENTION

The present invention provides a novel cytokine designated CD30-L, as well as isolated DNA encoding CD30-L protein, expression vectors comprising the isolated DNA, and a method for producing CD30-L by cultivating host cells containing the expression vectors under conditions appropriate for expression of the CD30-L protein. CD30-L is a ligand that binds to the Hodgkin's disease-associated antigen CD30 (a cell surface receptor). Antibodies directed against the CD30-L protein or an immunogenic

fragment thereof are also provided. Uses of CD30-L in diagnostic and therapeutic procedures are also disclosed.

DETAILED DESCRIPTION OF THE INVENTION

5 cDNA encoding a novel polypeptide that can act as a ligand for the Hodgkin's Disease-associated receptor known as CD30 has been isolated in accordance with the present invention. Also provided are expression vectors comprising the CD30 ligand (CD30-L) cDNA and methods for producing recombinant CD30-L polypeptides by cultivating host cells containing the expression vectors under conditions appropriate for
10 expression of CD30-L, and recovering the expressed CD30-L. Purified CD30-L protein is also encompassed by the present invention.

The present invention also provides CD30-L or antigenic fragments thereof that can act as immunogens to generate antibodies specific to the CD30-L immunogens. Monoclonal antibodies specific for CD30-L or antigenic fragments thereof thus can be
15 prepared.

The novel cytokine disclosed herein is a ligand for CD30, a receptor that is a member of the TNF/NGF receptor superfamily. Therefore, CD30-L is likely to be responsible for transducing a biological signal via CD30, which is known to be expressed on the surface of Hodgkin's Disease tumor cells.

20 One use of the CD30 ligand of the present invention is as a research tool for studying the pathogenesis of Hodgkin's Disease. As described in examples 8 and 13, CD30-L enhances the proliferation of the CD30⁺ neoplastic Hodgkin's Disease-derived lymphoma cell lines HDLM-2 and L-540, which are phenotypically T-cell-like. CD30-L did not produce a detectable effect on proliferation or viability of the B-cell-like,
25 CD30⁺ Hodgkin's Disease-derived lymphoma cell lines KM-H2 and L-428. The CD30-L of the present invention provides a means for investigating the roles that CD30-L and the cognate receptor may play in the etiology of Hodgkin's Disease.

CD30-L reduced proliferation of CD30⁺ large cell anaplastic lymphoma cell lines (one type of non-Hodgkin's lymphoma) (see examples 8 and 13). Thus, CD30-L
30 has potential use as a therapeutic agent. CD30-L also finds use in delivering diagnostic or therapeutic agents attached thereto to cells (e.g., malignant cells) that express the CD30 antigen.

The CD30 ligand also induces proliferation of T cells in the presence of an anti-CD3 co-stimulus. The CD30-L of the present invention thus is also useful as a
35 research tool for elucidating the roles that CD30 and CD30-L may play in the immune system. The inducible expression of CD30-L on normal T cells and macrophages, and

the presence of its receptor on activated T and B cells, is consistent with both autocrine and paracrine effects.

Upregulation of CD30 accompanying EBV, HTLV I and HTLV VII transformation also warrants further investigation, and the CD30-L provided herein is useful in such studies. HTLV I is the proximal cause of adult T cell Leukemia/Lymphoma. EBV has long been associated with Burkitt's lymphoma and nasopharyngeal carcinoma, and, overall, 50% of Hodgkin's lymphomas are EBV⁺ (reviewed in Klein, *Blood* 80:299 (1992)).

The CD30-L polypeptides of the present invention also may be employed in *in vitro* assays for detection of CD30 or CD30-L or the interactions thereof. Additional cell types expressing CD30 may be identified, for example.

The term "CD30-L" as used herein refers to a genus of polypeptides which are capable of binding CD30. Human CD30-L is within the scope of the present invention, as are CD30-L proteins derived from other mammalian species. As used herein, the term "CD30-L" includes membrane-bound proteins (comprising a cytoplasmic domain, a transmembrane region, and an extracellular domain) as well as truncated proteins that retain the CD30-binding property. Such truncated proteins include, for example, soluble CD30-L comprising only the extracellular (receptor binding) domain.

Isolation of a cDNA encoding murine CD30-L is described in examples 1-4 below. A human CD30-Fc fusion protein was prepared as described in example 1 for use in screening clones in a direct expression cloning procedure, to identify those expressing a protein that binds CD30.

Briefly, total RNA was isolated from a virally transformed human T-cell line designated HUT-102, which has been described by Durkop et al., *supra*, and Poiesz et al. (*PNAS USA* 77:7415-19, 1980). First strand cDNA was prepared using the total RNA as template. DNA encoding the extracellular domain of human CD30 was amplified by polymerase chain reaction (PCR) using primers based on the human CD30 sequence published by Durkop et al., *supra*, and the amplified DNA fragment was isolated. An expression vector comprising the CD30 extracellular domain DNA fused in-frame to the N-terminus of a human IgG1 Fc region DNA sequence was constructed and transfected into mammalian cells. The expressed protein was purified by a procedure that involved use of a protein G column (to which the Fc portion of the fusion protein binds).

Three activated murine helper T-cell lines were screened using a fluorescence activated cell sorting technique, and all three were found to bind a fluorescent derivative of the CD30-Fc protein. A cDNA library was prepared from one of the murine helper T-cell lines. cDNA from this library (in a mammalian expression vector that also

replicates in *E. coli*) was transfected into COS-7 (mammalian) cells, for isolation of clones expressing a CD30-binding protein by using a direct expression cloning technique. The clones were screened for ability to bind ^{125}I -CD30/Fc, and a positive clone was isolated. The recombinant vector isolated from the positive clone (murine
5 CD30-L cDNA in plasmid pDC202) was transformed into *E. coli* cells, deposited with the American Type Culture Collection on May 28, 1992, and assigned accession no. ATCC 69004. The deposit was made under the terms of the Budapest Treaty.

The murine CD30-L cDNA was radiolabeled and used as a probe to isolate human CD30-L cDNA by cross-species hybridization. Briefly, a cDNA library
10 prepared from activated human peripheral blood lymphocytes was screened with ^{32}P -labeled murine cDNA and a positive clone was isolated as described in Example 6. Human CD30-L DNA isolated from the positive clone was inserted into plasmid pGEMBL and then transformed into *E. coli* cells as described in Example 6. Samples of *E. coli* cells transformed with the recombinant vector were deposited with the
15 American Type Culture Collection on June 24, 1992, and assigned accession no. ATCC 69020. The deposit was made under the terms of the Budapest Treaty.

Additional murine and human CD30-L DNA sequences were isolated as described in example 7. The proteins encoded by the clones of example 7 comprise additional amino acids at the N-terminus, compared to the clones isolated in examples 4
20 and 6.

CD30-L proteins of the present invention thus include, but are not limited to, murine CD30-L proteins characterized by the N-terminal amino acid sequence Met-Gln-Val-Gln-Pro-Gly-Ser-Val-Ala-Ser-Pro-Trp (amino acids 1-12 of SEQ ID NO:19) or Met-Glu-Pro-Gly-Leu-Gln-Gln-Ala-Gly-Ser-Cys-Gly (amino acids 1-12 of SEQ ID
25 NO:6). Human CD30-L proteins characterized by the N-terminal amino acid sequence Met-His-Val-Pro-Ala-Gly-Ser-Val-Ala-Ser-His-Leu (amino acids 1-12 of SEQ ID NO:23) or Met-Asp-Pro-Gly-Leu-Gln-Gln-Ala-Leu-Asn-Gly-Met (amino acids 1-12 of SEQ ID NO:8) also are provided.

While a CD30/Fc fusion protein was employed in the screening procedure
30 described in example 4 below, labeled CD30 could be used to screen clones and candidate cell lines for expression of CD30-L proteins. The CD30/Fc fusion protein offers the advantage of being easily purified. In addition, disulfide bonds form between the Fc regions of two separate fusion protein chains, creating dimers. The dimeric CD30/Fc receptor was chosen for the potential advantage of higher affinity
35 binding of the CD30 ligand, in view of the possibility that the ligand being sought would be multimeric.

Further, other suitable fusion proteins comprising CD30 may be substituted for CD30/Fc in the screening procedures. Other fusion proteins can be made by fusing a DNA sequence for the ligand binding domain of CD30 to a DNA sequence encoding another polypeptide that is capable of affinity purification, for example, avidin or streptavidin. The resultant gene construct can be introduced into mammalian cells to express a fusion protein. Receptor/avidin fusion proteins can be purified by biotin affinity chromatography. The fusion protein can later be recovered from the column by eluting with a high salt solution or another appropriate buffer. Other antibody Fc regions may be substituted for the human IgG1 Fc region described in example 1. Other suitable Fc regions are defined as any region that can bind with high affinity to protein A or protein G, and include the Fc region of murine IgG1 or fragments of the human IgG1 Fc region, e.g., fragments comprising at least the hinge region so that interchain disulfide bonds will form.

cDNA encoding a CD30-L polypeptide may be isolated from other mammalian species by procedures analogous to those employed in isolating the murine CD30-L clone. For example, a cDNA library derived from a different mammalian species may be substituted for the murine cDNA library that was screened for binding of radioiodinated human CD30/Fc fusion protein in the direct expression cloning procedure described in example 4. Cell types from which cDNA libraries may be prepared may be chosen by the FACS selection procedure described in example 2, or any other suitable technique. As one alternative, mRNAs isolated from various cell lines can be screened by Northern hybridization to determine a suitable source of mammalian CD30-L mRNA for use in cloning a CD30-L gene.

Alternatively, one can utilize the murine or human CD30-L cDNAs described herein to screen cDNA derived from other mammalian sources for CD30-L cDNA using cross-species hybridization techniques. Briefly, an oligonucleotide based on the nucleotide sequence of the coding region (preferably the extracellular region) of the murine or human clone, or, preferably, the full length CD30-L cDNA, is prepared by standard techniques for use as a probe. The murine or human probe is used to screen a mammalian cDNA library or genomic library, generally under moderately stringent conditions.

CD30-L proteins of the present invention include, but are not limited to, murine CD30-L comprising amino acids 1-220 of SEQ ID NO:19 or 1-239 of SEQ ID NO:6; human CD30-L comprising amino acids 1-215 of SEQ ID NO:23 or 1-234 of SEQ ID NO:8; and proteins that comprise N-terminal, C-terminal, or internal truncations of the foregoing sequences, but retain the desired biological activity. Examples include murine CD30-L proteins comprising amino acids x to 239 of SEQ ID NO:6, wherein x

is 1-19 (i.e., the N-terminal amino acid is selected from amino acids 1-19 of SEQ ID NO:6, and the C-terminal amino acid is amino acid 239 of SEQ ID NO:6.) As described in example 7, amino acids 1-19 of the SEQ ID NO:6 sequence are not essential for binding of murine CD30-L to the CD30 receptor. Also provided by the present invention are human CD30-L proteins comprising amino acids y to 234 of SEQ ID NO:8 wherein y is 1-19 (i.e., the N-terminal amino acid is any one of amino acids 1-19 of SEQ ID NO:8, and amino acid 234 is the C-terminal amino acid. Such proteins, truncated at the N-terminus, are capable of binding CD30, as discussed in example 7.

One embodiment of the present invention provides soluble CD30-L polypeptides. Soluble CD30-L polypeptides comprise all or part of the extracellular domain of a native CD30-L but lack the transmembrane region that would cause retention of the polypeptide on a cell membrane. Since the CD30-L protein lacks a signal peptide, a heterologous signal peptide is fused to the N-terminus of a soluble CD30-L protein to promote secretion thereof, as described in more detail below. The signal peptide is cleaved from the CD30-L protein upon secretion from the host cell. The soluble CD30-L polypeptides that may be employed retain the ability to bind the CD30 receptor. Soluble CD30-L may also include part of the transmembrane region or part of the cytoplasmic domain or other sequences, provided that the soluble CD30-L protein is capable of being secreted.

Soluble CD30-L may be identified (and distinguished from its non-soluble membrane-bound counterparts) by separating intact cells which express the desired protein from the culture medium, e.g., by centrifugation, and assaying the medium (supernatant) for the presence of the desired protein. The culture medium may be assayed using procedures which are similar or identical to those described in the examples below. The presence of CD30-L in the medium indicates that the protein was secreted from the cells and thus is a soluble form of the desired protein.

The use of soluble forms of CD30-L is advantageous for certain applications. Purification of the proteins from recombinant host cells is facilitated, since the soluble proteins are secreted from the cells.

Examples of soluble CD30-L polypeptides include those comprising the entire extracellular domain of a native CD30-L protein or a fragment of said extracellular domain that is capable of binding CD30. One such soluble CD30-L comprises amino acids 49 (Gln) through 220 (Asp) of the murine CD30-L sequence of SEQ ID NO:19. Other soluble CD30-L polypeptides comprise amino acids z to 215 (Asp) of the human CD30-L sequence of SEQ ID NO:23, wherein z is 44, 45, 46, or 47. In other words, the N-terminal amino acid of the soluble human CD30-L is selected from the amino

acids in positions 44-47 of SEQ ID NO:23. DNA sequences encoding such soluble human CD30-L polypeptides include, but are not limited to, DNA sequences comprising a nucleotide sequence selected from the group consisting of nucleotides 130-645, 133-645, 136-645, and 139-645 of SEQ ID NO:22. Such sequences encode polypeptides comprising amino acids 44-215, 45-215, 46-215, and 47-215, respectively, of SEQ ID NO:23. Production of one such soluble human CD30-L protein, in the form of a fusion protein comprising amino acids 47-215 of SEQ ID NO:23 and an antibody Fc polypeptide, is illustrated in example 11.

Truncated CD30-L, including soluble polypeptides, may be prepared by any of a number of conventional techniques. In the case of recombinant proteins, a DNA fragment encoding a desired fragment may be subcloned into an expression vector. Alternatively, a desired DNA sequence may be chemically synthesized using known techniques. DNA fragments also may be produced by restriction endonuclease digestion of a full length cloned DNA sequence, and isolated by electrophoresis on agarose gels. Linkers containing restriction endonuclease cleavage site(s) may be employed to insert the desired DNA fragment into an expression vector, or the fragment may be digested at cleavage sites naturally present therein. The well known polymerase chain reaction procedure also may be employed to isolate a DNA sequence encoding a desired protein fragment.

In another approach, enzymatic treatment (e.g., using Bal 31 exonuclease) may be employed to delete terminal nucleotides from a DNA fragment to obtain a fragment having a particular desired terminus. Among the commercially available linkers are those that can be ligated to the blunt ends produced by Bal 31 digestion, and which contain restriction endonuclease cleavage site(s). Alternatively, oligonucleotides that reconstruct the N- or C-terminus of a DNA fragment to a desired point may be synthesized. The oligonucleotide may contain a restriction endonuclease cleavage site upstream of the desired coding sequence and position an initiation codon (ATG) at the N-terminus of the coding sequence.

The present invention provides purified CD30-L polypeptides, both recombinant and non-recombinant. Variants and derivatives of native CD30-L proteins that retain the desired biological activity are also within the scope of the present invention. CD30-L variants may be obtained by mutations of nucleotide sequences coding for native CD30-L polypeptides. A CD30-L variant, as referred to herein, is a polypeptide substantially homologous to a native CD30-L, but which has an amino acid sequence different from that of native CD30-L (human, murine or other mammalian species) because of one or a plurality of deletions, insertions or substitutions.

The variant amino acid sequence preferably is at least 80% identical to a native CD30-L amino acid sequence, most preferably at least 90% identical. The degree of homology (percent identity) may be determined, for example, by comparing sequence information using the GAP computer program, version 6.0 described by Devereux et al. (5) (*Nucl. Acids Res.* 12:387, 1984) and available from the University of Wisconsin Genetics Computer Group (UWGC). The GAP program utilizes the alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), as revised by Smith and Waterman (*Adv. Appl. Math* 2:482, 1981). The preferred default parameters for the GAP program include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for nucleotides, and the weighted comparison matrix of Gribskov and Burgess, (*Nucl. Acids Res.* 14:6745, 1986, as described by Schwartz and Dayhoff, eds., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358, 1979; (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

Alterations of the native amino acid sequence may be accomplished by any of a number of known techniques. Mutations can be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion.

Alternatively, oligonucleotide-directed site-specific mutagenesis procedures can be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Exemplary methods of making such alterations are disclosed by Walder et al. (*Gene* 42:133, 1986); Bauer et al. (*Gene* 37:73, 1985); Craik (*BioTechniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); and U.S. Patent Nos. 4,518,584 and 4,737,462, which are incorporated by reference herein.

Variants may comprise conservatively substituted sequences, meaning that a given amino acid residue is replaced by a residue having similar physiochemical characteristics. Examples of conservative substitutions include substitution of one aliphatic residue for another, such as Ile, Val, Leu, or Ala for one another, or substitutions of one polar residue for another, such as between Lys and Arg; Glu and Asp; or Gln and Asn. Other such conservative substitutions, for example, substitutions of entire regions having similar hydrophobicity characteristics, are well known.

CD30-L also may be modified to create CD30-L derivatives by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, lipids, phosphate, acetyl groups and the like. Covalent derivatives of CD30-L

may be prepared by linking the chemical moieties to functional groups on CD30-L amino acid side chains or at the N-terminus or C-terminus of a CD30-L polypeptide or the extracellular domain thereof. Other derivatives of CD30-L within the scope of this invention include covalent or aggregative conjugates of CD30-L or its fragments with
5 other proteins or polypeptides, such as by synthesis in recombinant culture as N-terminal or C-terminal fusions. For example, the conjugate may comprise a signal or leader polypeptide sequence (e.g. the α -factor leader of *Saccharomyces*) at the N-terminus of a soluble CD30-L polypeptide. The signal or leader peptide co-translationally or post-translationally directs transfer of the conjugate from its site of
10 synthesis to a site inside or outside of the cell membrane or cell wall.

CD30-L polypeptide fusions can comprise peptides added to facilitate purification and identification of CD30-L. Such peptides include, for example, poly-His or the antigenic identification peptides described in U.S. Patent No. 5,011,912 and in Hopp et al., *BioTechnology* 6:1204, 1988. One such peptide is the FLAG®
15 peptide, Asp-Tyr-Lys-Asp-Asp-Asp-Lys (DYKDDDDK) (SEQ ID NO:15), which is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody enabling rapid assay and facile purification of expressed recombinant protein. This sequence is also specifically cleaved by bovine mucosal enterokinase at the residue immediately following the Asp-Lys pairing. Fusion proteins capped with this peptide
20 may also be resistant to intracellular degradation in *E. coli*. A murine hybridoma designated 4E11 produces a monoclonal antibody that binds the peptide DYKDDDDK (SEQ ID NO:15) in the presence of certain divalent metal cations (as described in U.S. Patent 5,011,912) and has been deposited with the American Type Culture Collection under accession no HB 9259.

The present invention further includes CD30-L polypeptides with or without associated native-pattern glycosylation. CD30-L expressed in yeast or mammalian expression systems (e.g., COS-7 cells) may be similar to or significantly different from a native CD30-L polypeptide in molecular weight and glycosylation pattern, depending upon the choice of expression system. Expression of CD30-L polypeptides
30 in bacterial expression systems, such as *E. coli*, provides non-glycosylated molecules.

DNA constructs that encode various additions or substitutions of amino acid residues or sequences, or deletions of terminal or internal residues or sequences not needed for biological activity or binding can be prepared. For example, N-glycosylation sites in the CD30-L extracellular domain can be modified to preclude
35 glycosylation while allowing expression of a homogeneous, reduced carbohydrate analog using yeast or mammalian expression systems. N-glycosylation sites in eukaryotic polypeptides are characterized by an amino acid triplet Asn-X-Y, wherein X

is any amino acid except Pro and Y is Ser or Thr. Appropriate modifications to the nucleotide sequence encoding this triplet will result in substitutions, additions or deletions that prevent attachment of carbohydrate residues at the Asn side chain. Alteration of a single nucleotide, chosen so that Asn is replaced by a different amino acid, for example, is sufficient to inactivate an N-glycosylation site. Known procedures for inactivating N-glycosylation sites in proteins include those described in U.S. Patent 5,071,972 and EP 276,846.

In another example, sequences encoding Cys residues that are not essential for biological activity can be altered to cause the Cys residues to be deleted or replaced with other amino acids, preventing formation of incorrect intramolecular disulfide bridges upon renaturation. Other variants are prepared by modification of adjacent dibasic amino acid residues to enhance expression in yeast systems in which KEX2 protease activity is present. EP 212,914 discloses the use of site-specific mutagenesis to inactivate KEX2 protease processing sites in a protein. KEX2 protease processing sites are inactivated by deleting, adding or substituting residues to alter Arg-Arg, Arg-Lys, and Lys-Arg pairs to eliminate the occurrence of these adjacent basic residues. Lys-Lys pairings are considerably less susceptible to KEX2 cleavage, and conversion of Arg-Lys or Lys-Arg to Lys-Lys represents a conservative and preferred approach to inactivating KEX2 sites. The resulting muteins are less susceptible to cleavage by the KEX2 protease at locations other than the yeast α -factor leader sequence, where cleavage upon secretion is intended.

Naturally occurring CD30-L variants are also encompassed by the present invention. Examples of such variants are proteins that result from alternative mRNA splicing events (since CD30-L presumably is encoded by a multi-exon gene) or from proteolytic cleavage of the CD30-L protein, wherein the CD30-binding property is retained. Alternative splicing of mRNA may yield a truncated but biologically active CD30-L protein, such as a naturally occurring soluble form of the protein, for example. Variations attributable to proteolysis include, for example, differences in the N- or C-termini upon expression in different types of host cells, due to proteolytic removal of one or more terminal amino acids from the CD30-L protein (generally from 1-5 terminal amino acids).

Nucleic acid sequences within the scope of the present invention include isolated DNA and RNA sequences that hybridize to the CD30-L nucleotide sequences disclosed herein under conditions of moderate or severe stringency, and which encode biologically active CD30-L. Moderate stringency hybridization conditions refer to conditions described in, for example, Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2 ed. Vol. 1, pp. 1.101-104, Cold Spring Harbor Laboratory

Press, (1989). Conditions of moderate stringency, as defined by Sambrook et al., include use of a prewashing solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0) and hybridization conditions of about 55°C, 5 X SSC, overnight. Conditions of severe stringency include higher temperatures of hybridization and washing. The skilled
5 artisan will recognize that the temperature and wash solution salt concentration may be adjusted as necessary according to factors such as the length of the probe. One embodiment of the invention is directed to DNA sequences that will hybridize under severely stringent conditions to a DNA sequence comprising the coding region of a CD30-L clone disclosed herein. The severely stringent conditions include hybridization
10 at 68°C followed by washing in 0.1X SSC/0.1% SDS at 63-68°C.

The present invention thus provides isolated DNA sequences encoding biologically active CD30-L, selected from: (a) DNA derived from the coding region of a native mammalian CD30-L gene (e.g., DNA comprising the nucleotide sequence presented in SEQ ID NOS: 5, 7, 18, or 22; (b) DNA capable of hybridization to a
15 DNA of (a) under moderately (or severely) stringent conditions and which encodes biologically active CD30-L; and (c) DNA which is degenerate as a result of the genetic code to a DNA defined in (a) or (b) and which encodes biologically active CD30-L. CD30-L proteins encoded by the DNA sequences of (a), (b) and (c) are encompassed by the present invention.

20 Examples of CD30-L proteins encoded by DNA that varies from the native DNA sequences of SEQ ID NOS: 5, 7, 18, or 22, wherein the variant DNA will hybridize to a native DNA sequence under moderately stringent conditions, include, but are not limited to, CD30-L fragments (soluble or membrane-bound) and CD30-L proteins comprising inactivated N-glycosylation site(s), inactivated KEX2 protease
25 processing site(s), or conservative amino acid substitution(s), as described above. CD30-L proteins encoded by DNA derived from other mammalian species, wherein the DNA will hybridize to the human or murine DNA of SEQ ID NOS: 5, 7, 18, or 22, are also encompassed.

Variants possessing the requisite ability to bind CD30 may be identified by any
30 suitable assay. Biological activity of CD30-L may be determined, for example, by competition for binding to the ligand binding domain of CD30 (i.e. competitive binding assays).

One type of a competitive binding assay for CD30-L polypeptide uses a radiolabeled, soluble human or murine CD30-L and intact cells expressing cell surface
35 CD30 (e.g., cell lines such as HUT102, described by Durkop et al., *supra*). Instead of intact cells, one could substitute soluble CD30 bound to a solid phase (such as a CD30/Fc fusion protein bound to a Protein A or Protein G column through interaction

with the Fc region of the fusion protein). Another type of competitive binding assay utilizes radiolabeled soluble CD30 such as a CD30/Fc fusion protein, and intact cells expressing CD30-L. Alternatively, soluble CD30-L could be bound to a solid phase.

Competitive binding assays can be performed using standard methodology.

- 5 For example, radiolabeled murine CD30-L can be used to compete with a putative CD30-L homolog to assay for binding activity against surface-bound CD30. Qualitative results can be obtained by competitive autoradiographic plate binding assays, or Scatchard plots may be utilized to generate quantitative results.

- Competitive binding assays with intact cells expressing CD30 can be performed
10 by two methods. In a first method, cells expressing cell surface CD30 are grown either in suspension or by adherence to tissue culture plates. Adherent cells can be removed by treatment with 5 mM EDTA treatment for ten minutes at 37° C. In a second method, transfected COS cells expressing membrane-bound CD30 can be used. COS cells or another mammalian cell can be transfected with human CD30 cDNA in an appropriate
15 vector to express full length CD30 with an extracellular region.

- Alternatively, soluble CD30 can be bound to a solid phase such as a column chromatography matrix or a similar substrate suitable for analysis for the presence of a detectable moiety such as ¹²⁵I. Binding to a solid phase can be accomplished, for example, by obtaining a CD30/Fc fusion protein and binding it to a protein A or protein
20 G-containing matrix.

- Another means to measure the biological activity of CD30-L (including variants) is to utilize conjugated, soluble CD30 (for example, ¹²⁵I-CD30/Fc) in competition assays similar to those described above. In this case, however, intact cells expressing CD30-L, or soluble CD30-L bound to a solid substrate, are used to measure
25 competition for binding of labeled, soluble CD30 to CD30-L by a sample containing a putative CD30-L variant.

- The CD30-L of the present invention can be used in a binding assay to detect cells expressing CD30. For example, CD30-L or an extracellular domain or a fragment thereof can be conjugated to a detectable moiety such as ¹²⁵I. Radiolabeling with ¹²⁵I
30 can be performed by any of several standard methodologies that yield a functional ¹²⁵I-CD30-L molecule labeled to high specific activity. Alternatively, another detectable moiety such as an enzyme that can catalyze a colorimetric or fluorometric reaction, biotin or avidin may be used. Cells to be tested for CD30 expression can be contacted with conjugated CD30-L. After incubation, unbound conjugated CD30-L is removed
35 and binding is measured using the detectable moiety.

The CD30 ligand proteins disclosed herein also may be employed to measure the biological activity of CD30 protein in terms of binding affinity for CD30-L. To

illustrate, CD30-L may be employed in a binding affinity study to measure the biological activity of a CD30 protein that has been stored at different temperatures, or produced in different cell types. The biological activity of a CD30 protein thus can be ascertained before it is used in a research study, for example.

- 5 CD30-L proteins find use as reagents that may be employed by those conducting "quality assurance" studies, e.g., to monitor shelf life and stability of CD30 protein under different conditions. CD30 ligands may be used in determining whether biological activity is retained after modification of a CD30 protein (e.g., chemical modification, truncation, mutation, etc.). The binding affinity of the modified CD30
- 10 protein for a CD30-L is compared to that of an unmodified CD30 protein to detect any adverse impact of the modifications on biological activity of CD30.

- A different use of a CD30 ligand is as a reagent in protein purification procedures. CD30-L or CD30-L/Fc fusion proteins may be attached to a solid support material by conventional techniques and used to purify CD30 by affinity
- 15 chromatography.

- CD30-L polypeptides also find use as carriers for delivering agents attached thereto to cells bearing the CD30 cell surface antigen. As discussed above, CD30 has been detected on cells that include, but are not limited to, cells associated with various lymphoid malignancies, e.g., Hodgkin's Disease tumor cells and certain non-
- 20 Hodgkin's lymphoma cells, e.g., large cell anaplastic lymphoma (LCAL) cells. CD30⁺ LCALs are characterized by the presence of strong CD30 surface expression on the anaplastic lymphoma cells (Stein et al., *Blood* 66:848, 1985). CD30-L polypeptides thus can be used to deliver diagnostic or therapeutic agents to these cells (or to other cell types found to express CD30 on the cell surface) in *in vitro* or *in vivo* procedures.
- 25 CD30⁺ cells are contacted with a conjugate comprising a diagnostic or therapeutic agent attached to a CD30-L polypeptide. The CD30-L binds to the target cells, thus allowing detection thereof (in the case of diagnostic agents) or treatment thereof (with therapeutic agents).

- One example of such use is to expose a CD30⁺ lymphoma cell line to a
- 30 therapeutic agent/CD30-L conjugate to assess whether the agent exhibits cytotoxicity toward the lymphoma cells. A number of different therapeutic agents attached to CD30-L may be included in an assay to detect and compare the cytotoxic effects of the agents on the lymphoma cells. CD30-L/diagnostic agent conjugates may be employed to detect the presence of CD30⁺ cells *in vitro* or *in vivo*.

- 35 Diagnostic and therapeutic agents that may be attached to a CD30-L polypeptide include, but are not limited to, drugs, toxins, radionuclides, chromophores, enzymes that catalyze a colorimetric or fluorometric reaction, and the like, with the particular

agent being chosen according to the intended application. Examples of drugs include those used in treating various forms of cancer, e.g., mechlorethamine, procarbazine, prednisone, dacarbazine, nitrogen mustards such as L-phenylalanine nitrogen mustard or cyclophosphamide, intercalating agents such as cis-diaminodichloroplatinum, antimetabolites such as 5-fluorouracil, vinca alkaloids such as vincristine or vinblastine, and antibiotics such as calicheamycin, bleomycin, doxorubicin, daunorubicin, and derivatives thereof. Combinations of such drugs, attached to CD30-L, may be employed. Among the toxins are ricin, abrin, saporin toxin, diphtheria toxin, *Pseudomonas aeruginosa* exotoxin A, ribosomal inactivating proteins, mycotoxins such as trichothecenes, and derivatives and fragments (e.g., single chains) thereof. Radionuclides suitable for diagnostic use include, but are not limited to, ¹²³I, ¹³¹I, ^{99m}Tc, ¹¹¹In, and ⁷⁶Br. Radionuclides suitable for therapeutic use include, but are not limited to, ¹³¹I, ²¹¹At, ⁷⁷Br, ¹⁸⁶Re, ¹⁸⁸Re, ²¹²Pb, ²¹²Bi, ¹⁰⁹Pd, ⁶⁴Cu, and ⁶⁷Cu.

Such agents may be attached to the CD30-L by any suitable conventional procedure. CD30-L, being a protein, comprises functional groups on amino acid side chains that can be reacted with functional groups on a desired agent to form covalent bonds, for example. The agent may be covalently linked to CD30-L via an amide bond, hindered disulfide bond, acid-cleavable linkage, and the like, which are among the conventional linkages chosen according to such factors as the structure of the desired agent. Alternatively, CD30-L or the agent to be linked thereto may be derivatized to generate or attach a desired reactive functional group. The derivatization may involve attachment of one of the bifunctional coupling reagents available for linking various molecules to proteins (Pierce Chemical Company, Rockford, Illinois). A number of techniques for radiolabeling proteins are known. One such method involves use of the IODO-GEN reagent (Pierce Chemical Company) to radioiodinate a CD30-L polypeptide. Radionuclide metals may be attached to CD30-L by using a suitable bifunctional chelating agent, examples of which are described in U.S. patents 4,897,255 and 4,965,392.

Conjugates comprising CD30-L and a suitable diagnostic or therapeutic agent (preferably covalently linked) are thus prepared. The conjugates are administered or otherwise employed in an amount appropriate for the particular application.

Preferred therapeutic agents are radionuclides and drugs. In one embodiment of the invention, the anti-tumor drug calicheamycin is attached to a soluble human CD30 ligand polypeptide.

As illustrated in examples 8 and 13, CD30 ligand polypeptides of the present invention have been found to reduce proliferation of L₅₁₇₈Y cell lines. The CD30-L was employed in unlabeled form, i.e., did not have any therapeutic agent attached thereto.

Thus, one embodiment of the present invention is directed to a method for inhibiting proliferation of CD30⁺ LCAL cells by contacting said cells with a CD30-L polypeptide. The present invention further provides a method for treating LCAL, involving administering a therapeutically effective amount of a CD30-L polypeptide to a patient afflicted with LCAL.

Hybridoma cell lines that produce two monoclonal antibodies (MAbs) designated M44 and M67 were generated as described in example 12, using a soluble human CD30/Fc fusion protein as the immunogen. The M44 and M67 MAbs exhibited certain biological activities in common with CD30-L, one of which is reduction of proliferation of LCAL cells. Thus, the present invention also provides a method of inhibiting proliferation of CD30⁺ LCAL cells by contacting said cells with M44, M67, or a combination thereof. The M44 or M67 antibodies may be substituted for CD30-L in the above-described method for treating LCAL patients. M44 and M67 are also useful for delivering diagnostic or cytotoxic agents attached thereto to any CD30⁺ cells. "Humanized" or chimeric versions of these antibodies (e.g., comprising a human constant region), may be produced by known techniques and employed in the foregoing methods. Antigen-binding antibody fragments (e.g., Fab, Fab', or F(ab')₂ fragments) also may be employed.

Oligomeric Forms of CD30-L

CD30-L polypeptides may exist as oligomers, such as dimers or trimers. Oligomers may be linked by disulfide bonds formed between cysteine residues on different CD30-L polypeptides. In one embodiment of the invention, a CD30-L dimer is created by fusing CD30-L to the Fc region of an antibody (IgG1) in a manner that does not interfere with binding of CD30-L to the CD30 ligand binding domain. The polypeptide preferably is fused to the N-terminus of a soluble CD30-L (comprising only the extracellular domain). A procedure for isolating DNA encoding an IgG1 Fc region for use in preparing fusion proteins is presented in example 1 below. A gene fusion encoding the CD30-L/Fc fusion protein is inserted into an appropriate expression vector. The CD30-L/Fc fusion proteins are allowed to assemble much like antibody molecules, whereupon interchain disulfide bonds form between Fc polypeptides, yielding divalent CD30-L. If fusion proteins are made with both heavy and light chains of an antibody, it is possible to form a CD30-L oligomer with as many as four CD30-L extracellular regions.

Alternatively, one can link multiple copies of CD30-L *via* peptide linkers. A fusion protein comprising two or more copies of CD30-L (preferably soluble CD30-L polypeptides), separated by peptide linkers, may be produced by recombinant DNA

technology. Among the peptide linkers that may be employed are amino acid chains that are from 5 to 100 amino acids in length, preferably comprising amino acids selected from the group consisting of glycine, asparagine, serine, threonine, and alanine. The production of recombinant fusion proteins comprising peptide linkers is illustrated in United States Patent 5,073,627, for example, which is hereby incorporated by reference.

The present invention provides oligomers of CD30-L extracellular domains or fragments thereof, linked by disulfide bonds, or expressed as fusion proteins with or without spacer amino acid linking groups. For example, a dimer CD30-L molecule can be linked by an IgG Fc region linking group. Analysis of expressed recombinant CD30-L of the present invention by SDS-PAGE revealed both monomeric and oligomeric forms of the protein. The CD30-L proteins of the present invention are believed to form oligomers (disulfide-bonded dimers, trimers and higher oligomers) intracellularly. The oligomers then become attached to the cell surface via the transmembrane region of the protein.

Expression Systems

The present invention provides recombinant expression vectors for expression of CD30-L, and host cells transformed with the expression vectors. Any suitable expression system may be employed. The vectors include a CD30-L DNA sequence (e.g., a synthetic or cDNA-derived DNA sequence encoding a CD30-L polypeptide) operably linked to suitable transcriptional or translational regulatory nucleotide sequences, such as those derived from a mammalian, microbial, viral, or insect gene. Examples of regulatory sequences include transcriptional promoters, operators, or enhancers, an mRNA ribosomal binding site, and appropriate sequences which control transcription and translation initiation and termination. Nucleotide sequences are operably linked when the regulatory sequence functionally relates to the CD30-L DNA sequence. Thus, a promoter nucleotide sequence is operably linked to a CD30-L DNA sequence if the promoter nucleotide sequence controls the transcription of the CD30-L DNA sequence. The ability to replicate in the desired host cells, usually conferred by an origin of replication, and a selection gene by which transformants are identified, may additionally be incorporated into the expression vector.

In addition, sequences encoding appropriate signal peptides that are not native to the CD30-L gene can be incorporated into expression vectors. For example, a DNA sequence for a signal peptide (secretory leader) may be fused in frame to the CD30-L sequence so that the CD30-L is initially translated as a fusion protein comprising the signal peptide. A signal peptide fused to the N-terminus of a soluble CD30-L protein

promotes extracellular secretion of the CD30-L. The signal peptide is cleaved from the CD30-L polypeptide upon secretion of CD30-L from the cell. Signal peptides are chosen according to the intended host cells, and representative examples are described below.

- 5 Suitable host cells for expression of CD30-L polypeptides include prokaryotes, yeast or higher eukaryotic cells. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described, for example, in Pouwels et al. *Cloning Vectors: A Laboratory Manual*, Elsevier, New York, (1985). Cell-free translation systems could also be employed to produce CD30-L
10 polypeptides using RNAs derived from DNA constructs disclosed herein.

- Prokaryotes include gram negative or gram positive organisms, for example, *E. coli* or *Bacilli*. Suitable prokaryotic host cells for transformation include, for example, *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various other species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. In a prokaryotic host cell,
15 such as *E. coli*, a CD30-L polypeptide may include an N-terminal methionine residue to facilitate expression of the recombinant polypeptide in the prokaryotic host cell. The N-terminal Met may be cleaved from the expressed recombinant CD30-L polypeptide.

- Expression vectors for use in prokaryotic host cells generally comprise one or more phenotypic selectable marker genes. A phenotypic selectable marker gene is, for
20 example, a gene encoding a protein that confers antibiotic resistance or that supplies an autotrophic requirement. Examples of useful expression vectors for prokaryotic host cells include those derived from commercially available plasmids such as the cloning vector pBR322 (ATCC 37017). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells. An
25 appropriate promoter and a CD30-L DNA sequence are inserted into the pBR322 vector. Other commercially available vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, WI, USA).

- Promoter sequences commonly used for recombinant prokaryotic host cell
30 expression vectors include β -lactamase (penicillinase), lactose promoter system (Chang et al., *Nature* 275:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), tryptophan (trp) promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980; and EP-A-36776) and tac promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p. 412, 1982). A particularly useful prokaryotic host cell
35 expression system employs a phage λ P_L promoter and a ci857s thermolabile repressor sequence. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the λ P_L promoter include plasmid pHUB2 (resident in *E.*

coli strain JMB9 (ATCC 37092)) and pPLc28 (resident in *E. coli* RR1 (ATCC 53082)).

CD30-L alternatively may be expressed in yeast host cells, preferably from the *Saccharomyces* genus (e.g., *S. cerevisiae*). Other genera of yeast, such as *Pichia* or *Kluyveromyces*, may also be employed. Yeast vectors will often contain an origin of replication sequence from a 2 μ yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Suitable promoter sequences for yeast vectors include, among others, promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* 255:2073, 1980) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968; and Holland et al., *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Other suitable vectors and promoters for use in yeast expression are further described in Hitzeman, EPA-73,657. Another alternative is the glucose-repressible ADH2 promoter described by Russell et al. (*J. Biol. Chem.* 258:2674, 1982) and Beier et al. (*Nature* 300:724, 1982). Shuttle vectors replicable in both yeast and *E. coli* may be constructed by inserting DNA sequences from pBR322 for selection and replication in *E. coli* (Amp^r gene and origin of replication) into the above-described yeast vectors.

The yeast α -factor leader sequence may be employed to direct secretion of the CD30-L polypeptide. The α -factor leader sequence is often inserted between the promoter sequence and the structural gene sequence. See, e.g., Kurjan et al., *Cell* 30:933, 1982; Bitter et al., *Proc. Natl. Acad. Sci. USA* 81:5330, 1984; U.S. Patent 4,546,082; and EP 324,274. Other leader sequences suitable for facilitating secretion of recombinant polypeptides from yeast hosts are known to those of skill in the art. A leader sequence may be modified near its 3' end to contain one or more restriction sites. This will facilitate fusion of the leader sequence to the structural gene.

Yeast transformation protocols are known to those of skill in the art. One such protocol is described by Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929, 1978. The Hinnen et al. protocol selects for Trp⁺ transformants in a selective medium, wherein the selective medium consists of 0.67% yeast nitrogen base, 0.5% casamino acids, 2% glucose, 10 μ g/ml adenine and 20 μ g/ml uracil.

Yeast host cells transformed by vectors containing ADH2 promoter sequence may be grown for inducing expression in a "rich" medium. An example of a rich medium is one consisting of 1% yeast extract, 2% peptone, and 1% glucose

supplemented with 80 µg/ml adenine and 80 µg/ml uracil. Derepression of the ADH2 promoter occurs when glucose is exhausted from the medium.

Mammalian or insect host cell culture systems could also be employed to express recombinant CD30-L polypeptides. Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow and Summers, *BioTechnology* 6:47 (1988). Established cell lines of mammalian origin also may be employed. Examples of suitable mammalian host cell lines include the COS-7 line of monkey kidney cells (ATCC CRL 1651) (Gluzman et al., *Cell* 23:175, 1981), L cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells, HeLa cells, and BHK (ATCC CRL 10) cell lines, and the CV1/EBNA cell line derived from the African green monkey kidney cell line CV1 (ATCC CCL 70) as described by McMahan et al. (*EMBO J.* 10: 2821, 1991).

Transcriptional and translational control sequences for mammalian host cell expression vectors may be excised from viral genomes. Commonly used promoter sequences and enhancer sequences are derived from Polyoma virus, Adenovirus 2, Simian Virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide other genetic elements for expression of a structural gene sequence in a mammalian host cell. Viral early and late promoters are particularly useful because both are easily obtained from a viral genome as a fragment which may also contain a viral origin of replication (Fiers et al., *Nature* 273:113, 1978). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the *Hind* III site toward the *Bgl* I site located in the SV40 viral origin of replication site is included.

Expression vectors for use in mammalian host cells can be constructed as disclosed by Okayama and Berg (*Mol. Cell. Biol.* 3:280, 1983). A useful system for stable high level expression of mammalian cDNAs in C127 murine mammary epithelial cells can be constructed substantially as described by Cosman et al. (*Mol. Immunol.* 23:935, 1986). A useful high expression vector, PMLSV N1/N4, described by Cosman et al., *Nature* 312:768, 1984 has been deposited as ATCC 39890. Additional useful mammalian expression vectors are described in EP-A-0367566, and PCT Application WO 91/18982, incorporated by reference herein. The vectors may be derived from retroviruses. To achieve secretion of CD30-L (a type II protein lacking a native signal sequence), a heterologous signal sequence may be added. Examples of signal peptides useful in mammalian expression systems are the signal sequence for interleukin-7 (IL-7) described in United States Patent 4,965,195; the signal sequence for interleukin-2 receptor described in Cosman et al., *Nature* 312:768 (1984); the

interleukin-4 signal peptide described in EP 367,566; the type I interleukin-1 receptor signal peptide described in U.S. Patent 4,968,607; and the type II interleukin-1 receptor signal peptide described in EP 460,846. Each of these references describing signal peptides is hereby incorporated by reference.

5

CD30 Ligand Protein

The present invention provides substantially homogeneous CD30-L protein, which may be produced by recombinant expression systems as described above or purified from naturally occurring cells. The CD30-L is purified to substantial
10 homogeneity, as indicated by a single protein band upon analysis by SDS-polyacrylamide gel electrophoresis (SDS-PAGE).

In one embodiment of the present invention, CD30-L is purified from a cellular source using any suitable protein purification technique. The cells may, for example, be activated T-lymphocytes from a mammalian species of interest, such as the murine
15 cell line 7B9 described in examples 2 and 3 or induced human peripheral blood T-cells.

An alternative process for producing the CD30-L protein comprises culturing a host cell transformed with an expression vector comprising a DNA sequence that encodes CD30-L under conditions such that CD30-L is expressed. The CD30-L protein is then recovered from culture medium or cell extracts, depending upon the
20 expression system employed. As the skilled artisan will recognize, procedures for purifying the recombinant CD30-L will vary according to such factors as the type of host cells employed and whether or not the CD-30-L is secreted into the culture medium.

For example, when expression systems that secrete the recombinant protein are
25 employed, the culture medium first may be concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a purification matrix such as a gel filtration medium. Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl
30 (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfoethyl or carboxymethyl groups. Sulfoethyl groups are preferred. Finally, one or more reversed-phase high performance liquid chromatography (RP-
35 HPLC) steps employing hydrophobic RP-HPLC media, (e.g., silica gel having pendant methyl or other aliphatic groups) can be employed to further purify CD30-L.

Some or all of the foregoing purification steps, in various combinations, can be employed to provide a substantially homogeneous recombinant protein.

It is also possible to utilize an affinity column comprising the ligand binding domain of CD30 to affinity-purify expressed CD30-L polypeptides. CD30-L polypeptides can be removed from an affinity column in a high salt elution buffer and then dialyzed into a lower salt buffer for use. Alternatively, the affinity column may comprise an antibody that binds CD30-L. Example 5 describes a procedure for employing the CD30-L protein of the present invention to generate monoclonal antibodies directed against CD30-L.

Recombinant protein produced in bacterial culture is usually isolated by initial disruption of the host cells, centrifugation, extraction from cell pellets if an insoluble polypeptide, or from the supernatant fluid if a soluble polypeptide, followed by one or more concentration, salting-out, ion exchange, affinity purification or size exclusion chromatography steps. Finally, RP-HPLC can be employed for final purification steps. Microbial cells can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Transformed yeast host cells are preferably employed to express CD30-L as a secreted polypeptide. This simplifies purification. Secreted recombinant polypeptide from a yeast host cell fermentation can be purified by methods analogous to those disclosed by Urdal et al. (*J. Chromatog.* 296:171, 1984). Urdal et al. describe two sequential, reversed-phase HPLC steps for purification of recombinant human IL-2 on a preparative HPLC column.

The present invention provides pharmaceutical compositions comprising an effective amount of a purified CD30-L polypeptide and a suitable diluent, excipient, or carrier. Such carriers will be nontoxic to patients at the dosages and concentrations employed. Ordinarily, the preparation of such compositions entails combining a mammalian CD30-L polypeptide or derivative thereof with buffers, antioxidants such as ascorbic acid, low molecular weight (less than about 10 residues) peptides, proteins, amino acids, carbohydrates including glucose, sucrose, or dextrans, chelating agents such as EDTA, glutathione, or other stabilizers and excipients. Neutral buffered saline is one appropriate diluent.

For therapeutic use, the compositions are administered in a manner and dosage appropriate to the indication and the patient. As will be understood by one skilled in the pertinent field, a therapeutically effective dosage will vary according to such factors as the nature and severity of the disorder to be treated and the age, condition, and size of the patient. Administration may be by any suitable route, including but not limited to

intravenous injection, continuous infusion, local infusion during surgery, or sustained release from implants (gels, membranes, and the like).

- The compositions of the present invention may contain a CD30-L protein in any form described above, including variants, derivatives, biologically active fragments, and oligomeric forms thereof. CD30-L derived from the same mammalian species as the patient is generally preferred for use in pharmaceutical compositions. In one embodiment of the invention, the composition comprises a soluble human CD30-L protein. Such protein may be in the form of dimers comprising the extracellular domain of human CD30-L fused to an Fc polypeptide, as described above. In another embodiment of the invention, the pharmaceutical composition comprises a CD30-L polypeptide having a diagnostic or therapeutic agent attached thereto. Such compositions may be administered to diagnose or treat conditions characterized by CD30⁺ cells, e.g., Hodgkin's Disease or large cell anaplastic lymphomas, as discussed above. A composition comprising unlabeled CD30-L may be used in treating LCAL.
- 15 The foregoing compositions may additionally contain, or be co-administered with, additional agents effective in treating malignancies characterized by CD30⁺ cells.

Nucleic Acid Fragments

- The present invention further provides fragments of the CD30-L nucleotide sequences presented herein. Such fragments desirably comprise at least about 14 nucleotides of the sequence presented in SEQ ID NO:5 or SEQ ID NO:7. DNA and RNA complements of said fragments are provided herein, along with both single-stranded and double-stranded forms of the CD30-L DNA

- Among the uses of such CD30-L nucleic acid fragments is use as a probe.
- 25 Such probes may be employed in cross-species hybridization procedures to isolate CD30-L DNA from additional mammalian species. As one example, a probe corresponding to the extracellular domain of CD30-L may be employed. The probes also find use in detecting the presence of CD30-L nucleic acids in *in vitro* assays and in such procedures as Northern and Southern blots. Cell types expressing CD30-L can be identified. Such procedures are well known, and the skilled artisan can choose a probe of suitable length, depending on the particular intended application.

- Other useful fragments of the CD30-L nucleic acids are antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target CD30-L mRNA (sense) or CD30-L DNA (antisense) sequences. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of CD30-L cDNA. Such a fragment generally comprises at least about 14 nucleotides, preferably from about 14 to about 30

nucleotides. The ability to create an antisense or a sense oligonucleotide, based upon a cDNA sequence for a given protein is described in, for example, Stein and Cohen, *Cancer Res.* 48:2659, 1988 and van der Krol et al., *BioTechniques* 6:958, 1988.

Binding of antisense or sense oligonucleotides to target nucleic acid sequences results in the formation of duplexes that block translation (RNA) or transcription (DNA) by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. The antisense oligonucleotides thus may be used to block expression of CD30-L proteins.

Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO 91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable *in vivo* (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences. Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10448, and other moieties that increases affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence. Antisense or sense oligonucleotides may be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example, CaPO₄-mediated DNA transfection, electroporation, or other gene transfer vectors such as Epstein-Barr virus. Antisense or sense oligonucleotides are preferably introduced into a cell containing the target nucleic acid sequence by insertion of the antisense or sense oligonucleotide into a suitable retroviral vector, then contacting the cell with the retrovirus vector containing the inserted sequence, either *in vivo* or *ex vivo*. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see PCT Application US 90/02656).

Sense or antisense oligonucleotides may also be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand

binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

The following examples are provided to illustrate particular embodiments and not to limit the scope of the invention.

10 EXAMPLE 1: Preparation of Soluble CD30/Fc Fusion Protein

This example describes construction of a CD30/Fc-encoding vector to express a soluble CD30/Fc fusion protein for use in detecting cDNA clones encoding a CD30 ligand. A cDNA fragment encoding the extracellular region (ligand binding domain) of the CD30 human receptor was obtained using polymerase chain reaction (PCR) techniques, and is based upon the sequence published by Durkop et al. (*Cell* 68:421, 15 1992). The CD30 nucleotide sequence reported in Durkop et al. *supra* is presented in SEQ ID NO:1, and the amino acid sequence encoded thereby is presented in SEQ ID NO:2. The signal sequence comprises amino acids 1-18, and the transmembrane region comprises amino acids 391-407, of SEQ ID NO:2.

20 The CD30 cDNA used as a template in the PCR reaction was prepared as follows. Total RNA was isolated from a virally transformed human T-cell line designated HUT 102E. This cell line was derived by transforming T-cells with human T-cell lymphotropic virus 1 (HTLV-1) as described by Poiesz et al. (*PNAS USA* 77:7415-19, 1980). First strand cDNA was prepared using a SuperScript™ cDNA 25 synthesis kit available from GIBCO/BRL (Gaithersburg, Maryland). The resulting single-stranded cDNA was employed as the template in a PCR reaction.

The 5' primer employed in the PCR reaction was a single-stranded oligonucleotide (39-mer) of the sequence:

5' ATAGCGGCCGCCACCATGCGCGTCCTCCTCGCCGCGCTG 3'

30 This primer (SEQ ID NO:9) comprises a recognition site for the restriction endonuclease *NotI* (underlined) upstream of a sequence (double underline) encoding the first (N-terminal) eight amino acids of the CD30 sequence shown in SEQ ID NO:1, from methionine (encoded by the translation initiation codon ATG) through leucine at position eight.

35 The 3' primer employed in the PCR reaction was a single-stranded oligonucleotide (39-mer) of the sequence:

3' CAGCGAGAGAGGAGGTGCCCTTCTCCTCGGGTCTAGAACA 5'

This primer (SEQ ID NO:10) comprises a sequence (double underline) that is complementary to the sequence that encodes the last eight amino acids of the CD30 extracellular domain, i.e., amino acids 372 (Val) through 379 (Lys) shown in SEQ ID NO:1. The sequence CTCGGG that follows the CD30 sequence is complementary to codons for Glu and Pro. Glu and Pro are the first two amino acids of an antibody Fc fragment that is fused to the C-terminus of the CD30 fragment as described below. The primer also positions a recognition site for the restriction endonuclease *Bgl*II (underlined) downstream, for use in attaching a DNA sequence encoding the remainder of the Fc-encoding gene.

The PCR reaction may be conducted using any suitable procedure, such as those described in Sarki et al., *Science* 239:487 (1988); in *Recombinant DNA Methodology*, Wu et al., eds., Academic Press Inc., San Diego (1989), pp. 189-196; and in *PCR Protocols: A Guide to Methods and Applications*, Innis et al., eds., Academic Press, Inc. (1990). An example of a suitable PCR procedure is as follows.

All temperatures are in degrees centigrade. The following PCR reagents are added to a 0.5 ml Eppendorf microfuge tube: 10 μ l of 10X PCR buffer (500 mM KCl, 100 mM Tris-HCl, pH 8.3 at 25°C, 25 mM MgCl₂, and 1 mg/ml gelatin) (Perkins-Elmer Cetus, Norwalk, CN), 8 μ l of a 2.5 mM solution containing each dNTP (2 mM dATP, 2mM dCTP, 2mM dGTP and 2 mM dTTP), 2.5 units (0.5 μ l of standard 5000 units/ml solution) of *Taq* DNA polymerase (Perkins-Elmer Cetus), 1 ng of template DNA, 100 picomoles of each of the oligonucleotide primers, and water to a final volume of 100 μ l. The final mixture is then overlaid with 100 μ l paraffin oil. PCR is carried out using a DNA thermal cycler (Ericomp, San Diego, CA).

In a preferred procedure, the template was denatured at 94° for 5 minutes, followed by 5 cycles of 94° for 1 minute (denaturation), 48° for 1 min. (annealing), and 72° for 1 min. (extension); followed by 30 cycles of 94° for 1 min., 68° for 1 min., and 72° for 1 min., with the last cycle being followed by a final extension at 72° for 5 mins. An aliquot of the products of this PCR reaction was reamplified in a second PCR reaction, using the same conditions.

The desired DNA fragment amplified by this PCR reaction comprised a *Nor*I site upstream of a sequence encoding the entire extracellular domain of CD30, followed by a *Bgl*II site. The PCR reaction products were digested with *Nor*I and *Bgl*II, and the desired fragment was purified by gel electrophoresis.

A DNA sequence encoding an antibody Fc fragment, to be fused to the CD30-encoding DNA fragment, was prepared as follows. DNA encoding a single chain polypeptide derived from the Fc region of a human IgG1 antibody has been cloned into the *Spe*I site of the pBLUESCRIPT SK® vector, which is available from Stratagene

Cloning Systems, La Jolla, California. This plasmid vector is replicable in *E. coli* and contains a polylinker segment that includes 21 unique restriction sites. The DNA and encoded amino acid sequences of the cloned Fc cDNA coding region are presented in Figure 2. A unique *Bgl*III site has been introduced near the 5' end of the inserted Fc encoding sequence. Nucleotides 7-12 of SEQ ID NO:1 constitute the *Bgl*III recognition sequence.

The Fc polypeptide encoded by the DNA extends from the N-terminal hinge region to the native C-terminus, i.e., is an essentially full-length antibody Fc region. Fragments of Fc regions, e.g., those that are truncated at the C-terminal end, also may be employed. The fragments preferably contain multiple cysteine residues (at least the cysteine residues in the hinge reaction) to permit interchain disulfide bonds to form between the Fc polypeptide portions of two separate CD30/Fc fusion proteins, forming dimers as discussed above.

The recombinant vector containing the Fc sequence is digested with *Bgl*III (which cleaves only at the site shown in Figure 2) and *Not*I (which cleaves the vector in the multiple cloning site downstream of the Fc cDNA insert. The Fc-encoding fragment (about 720 bp in length) was isolated by conventional procedures using LMT agarose gel electrophoresis.

The *Not*I/*Bgl*III CD30-encoding DNA fragment and the *Bgl*III/*Not*I Fc-encoding DNA fragment prepared above were ligated into an expression vector designated pDC406 as follows. Plasmid pDC406, which has been described by McMahan et al. (*EMBO J.* 10:2821, 1991), is an expression vector for use in mammalian cells, but is also replicable in *E. coli* cells.

pDC406 contains origins of replication derived from SV40, Epstein-Barr virus and pBR322 and is a derivative of HAV-EO described by Dower et al., *J. Immunol.* 142:4314 (1989). pDC406 differs from HAV-EO by the deletion of the intron present in the adenovirus 2 tripartite leader sequence in HAV-EO. pDC406 was digested with *Not*I, which cleaves the plasmid in a multiple cloning site just 3' of the *Sal*I site, then treated with calf intestine alkaline phosphatase (CIAP) to prevent self ligation.

A three-way ligation to join the vector, Fc, and CD30 DNA fragments was conducted under conventional conditions, and *E. coli* cells were transformed with the ligation mixture. A plasmid of the desired size that was recovered from the *E. coli* cells was found to comprise the CD30/Fc gene fusion insert, but in the wrong orientation for expression. The CD30/Fc gene fusion was excised from this recombinant plasmid by *Not*I digestion and ligated to *Not*I-digested and CIAP-treated pDC406. *E. coli* cells were transformed with the ligation mixture. A recombinant plasmid containing the

insert in the desired orientation was isolated. The CD30 sequence was fused (in the same reading frame) to the downstream Fc sequence.

CD30/Fc fusion molecules preferably are synthesized in recombinant mammalian cell culture because they are generally too large and complex to be synthesized by prokaryotic expression methods. Examples of suitable mammalian cells for expressing a receptor/Fc fusion protein include CV-1 cells (ATCC CCL 70) and COS-7 cells (ATCC CRL 1651), both derived from monkey kidney.

The DNA construct pDC406/CD30/Fc was transfected into the monkey kidney cell line CV-1/EBNA (ATCC CRL 10478). In mammalian host cells such as CV1/EBNA, the CD30/Fc fusion protein is expressed off the HIV transactivating region (TAR) promoter. The CV-1/EBNA cell line was derived by transfection of the CV-1 cell line (ATCC CCL 70) with a gene encoding Epstein-Barr virus nuclear antigen-1 (EBNA-1) that constitutively expresses EBNA-1 driven from the human CMV intermediate-early enhancer/promoter as described by McMahan et al., *supra*. The EBNA-1 gene allows for episomal replication of expression vectors, such as pDC406, that contain the EBV origin of replication.

CV1-EBNA cells transfected with the pDC406/CD30/Fc vector were cultivated in roller bottles to allow transient expression of the fusion protein, which is secreted into the culture medium via the CD30 signal peptide. The CD30/Fc fusion protein was purified by affinity chromatography. Briefly, one liter of culture supernatant containing the CD30/Fc fusion protein was purified by filtering the supernatants (e.g., in a 0.45 μ filter) and applying the filtrate to a protein G affinity column (Schleicher and Schuell, Keene, NH) according to manufacturer's instructions. The Fc portion of the fusion protein is bound by the Protein G on the column. Bound fusion protein was eluted from the column and the purity confirmed on a silver stained SDS gel.

EXAMPLE 2: Screening of Cell Lines for Binding of CD30

This example describes screening of certain cell lines for the ability to bind a CD30/Fc fusion protein. Those cell lines found to be capable of binding CD30/Fc were considered to be candidates for use as nucleic acid sources in the attempt to clone CD30-L.

Biotinylation of CD30/Fc Fusion Proteins

The purified CD30/Fc fusion protein prepared in Example 1 was labeled with biotin for use in screening cell lines. CD30/Fc or control human IL-4R/Fc were biotinylated as follows: 50 μ g protein (200-500 μ g/ml in 0.1M NaHCO₃ pH 8.3) was incubated with 2 μ g (1 mg/ml in DMSO) Biotin-X-N-hydroxysuccinimide (NHS,

- Calbiochem, La Jolla, CA) for 30 min at room temperature. At the end of the incubation period, the reaction mixture was microfuged through a 1 ml Sephadex G-25 (Pharmacia) desalting column and the eluate adjusted to 100 µg/ml in PBS plus 0.02% NaN₃. Protein concentration of biotinylated CD30/Fc and hIL-4R/Fc was determined by micro-BCA assay (Pierce, Rockford, IL) with ultrapure bovine serum albumin as standard.

Flow cytometric staining with biotinylated Fc fusion proteins

- Cell lines such as those identified below are screened for binding of biotinylated CD30/Fc by the following procedure. Staining of 1×10^6 cells was carried out in round-bottomed 96-well microtiter plates in a volume of 20 µl. Cells were pre-incubated for 30 min at 4°C with 50 µl blocking solution consisting of 100 µg/ml human IgG1 + 2% goat serum in PBS + azide to prevent non-specific binding of labeled fusion proteins to Fc receptors. 150 µl PBS + azide was then added to the wells and cells were pelleted by centrifugation for 4 min at 1200 rpm. Pellets were resuspended in 20 µl of 5 µg/ml biotinylated CD30/Fc or biotinylated hIL-4R/Fc (as a specificity control) diluted in blocking solution. After 30-45 min incubation at 4°C, cells were washed X2 in PBS + azide and resuspended in 20 µl streptavidin-phycoerythrin (Becton Dickinson) diluted 1:5 in PBS + azide. After an additional 30 min, cells are washed x2 and are ready for analysis. If necessary, stained cells can be fixed in 1% formaldehyde, 1% fetal bovine serum in PBS + azide and stored at 4°C in the dark for analysis at a later time.

- Streptavidin binds to the biotin molecule which was attached to the CD30/Fc protein. Phycoerythrin is a fluorescent phycobiliprotein which serves as a detectable label. The level of fluorescence signal was then measured for each cell type using a FACScan® flow cytometer (Becton Dickinson).

Cell Lines to be Screened for CD30/Fc Binding

- Sheep red blood cell (SRBC)-specific helper T-cell lines designated 7C2 (TH1), 7B9 (TH0) and SBE11 (TH2) were derived by limiting dilution from primary antigen-induced cultures of murine C57BL/6 spleen cells. TH phenotypes of these clones were determined by their ability to secrete IL-2 and/or IL-4 in response to stimulation with the mitogen concanavalin A (ConA).

- Human peripheral blood T-cells were stimulated for 16 hours with 10 µg/ml of an anti-CD3 monoclonal antibody immobilized on plastic, prior to assay for CD30/Fc binding. The anti-CD3 MAb stimulates the T-cells through the CD3-T-cell receptor (TCR) complex.

Biotinylated CD30/Fc binding

Murine T-cell lines 7C2, 7B9 and SBE11 showed significant binding of biotinylated CD30/Fc over that seen with control IL-4R/Fc, after stimulation for 18 hours with 3 µg/ml Con A. 7C2 cells were also assayed after 6 hours stimulation with Con A, and specific binding of labeled CD30/Fc was seen. The anti-CD3 MAb activated human T-cells showed significant binding of biotinylated CD30/Fc. Binding of biotinylated CD30/Fc was not detected on any of these cell lines in the absence of stimulation.

Any of the cell lines that demonstrated binding of CD30/Fc may be used as a source of nucleic acid in an attempt to isolate a CD30-L-encoding DNA sequence. A cDNA library may be prepared from any of the three Con A stimulated murine T-cell lines or the activated human peripheral blood T-cells, and screened to identify CD30-L cDNA using the direct expression cloning strategy described below, for example. Other types of activated T-cells may be screened for CD30 binding to identify additional suitable nucleic acid sources. The cells may be derived from human, murine, or other mammalian sources, including but not limited to rat, bovine, porcine, or various primate cells. Further, the T-cells may be stimulated with mitogens other than ConA or otherwise activated by conventional techniques. It is to be noted that human CD30/Fc was successfully employed to screen both human and murine cell lines in the foregoing assay (i.e., human CD30/Fc binds to a ligand on both the human and the murine cell lines tested).

EXAMPLE 3: Preparation of cDNA Library Derived from Activated Murine Helper T-cells

This example describes preparation of a cDNA library for expression cloning of murine CD30-L. The library was prepared from the murine helper T-cell line designated 7B9 (described above and in Mosley et al., *Cell* 59:335, 1989), which was stimulated for 6 hours with 3 µg/ml Con A. The library construction technique was substantially similar to that described by Ausubel et al., eds., *Current Protocols In Molecular Biology*, Vol. 1, (1987). Briefly, total RNA was extracted from the 7B9 cell line and poly (A)⁺ mRNA was isolated by oligo dT cellulose chromatography. Double-stranded cDNA was made substantially as described by Gubler et al., *Gene* 25:263, 1983. Poly(A)⁺ mRNA fragments were converted to RNA-cDNA hybrids by reverse transcriptase using random hexanucleotides as primers. The RNA-cDNA hybrids were then converted into double-stranded cDNA fragments using RNAase H in combination with DNA polymerase I. The resulting double-stranded cDNA was blunt-ended with T4 DNA polymerase.

Unkinased (i.e. unphosphorylated) *Bgl*III adaptors:

5'- GATCTGGCAACGAAGGTACCATGG -3' (SEQ ID NO:11)

ACCGTTGCTTCCATGGTACC -5' (SEQ ID NO:12)

- were ligated to 5' ends of the resulting blunt-ended cDNA, using the adaptor cloning method described in Haymerle et al., *Nucleic Acids Res.* 14:8615, 1986. Only the 24-mer oligonucleotide (top strand) will covalently bond to the cDNA during the ligation reaction. Non-covalently bound adaptors (including the 20-mer oligonucleotide above) were removed by gel filtration chromatography at 68°C. This left 24 nucleotide non-self-complementary overhangs on cDNA. The cDNA was inserted into pDC202, a mammalian expression vector that also replicates in *E. coli*. pDC202 is derived from pDC201 (Sims et al., *Nature* 241:585, 1988). The plasmid pDC201 was assembled from (i) the SV40 origin of replication, enhancer, and early and late promoters; (ii) the adenovirus-2 major late promoter and tripartite leader; (iii) SV40 polyadenylation and transcription termination signals; (iv) adenovirus-2 virus-associated RNA genes (VAI and VAII); and (v) pMSLV (Cosman et al., *Nature* 312:768, 1984). The multiple cloning site contains recognition sites for *Kpn* I, *Sma* I, and *Bgl* II. Certain extraneous vector sequences bordering the VA genes were excised from pDC201 to create pDC202. Each of the above-named features of pDC201 is present in pDC202 as well. pDC202 was digested with *Bgl*III and *Bgl* II adaptors were ligated thereto as described for the cDNA above, except that the bottom strand of the adaptor (the 20-mer) is covalently bound to the vector, rather than the 24-mer ligated to the cDNA. A single-stranded extension complementary to that added to the cDNA thus was added to the *Bgl*III-digested vector. The 5' ends of the adapted vector and cDNA were phosphorylated and the two DNA species were then ligated in the presence of T4 polynucleotide kinase. Dialyzed ligation mixtures were electroporated into *E. coli* strain DH5 α and transformants selected on ampicillin plates.

- To create an expression cloning library, the recombinant vectors containing 7B9-derived cDNA were transferred from *E. coli* to mammalian host cells. Plasmid DNA was isolated from pools of transformed *E. coli* and transfected into a sub-confluent layer of COS-7 cells using standard techniques. The transfected cells were cultured for two to three days on chambered glass slides (Lab-Tek) to permit transient expression of the inserted DNA sequences.

EXAMPLE 4: Isolation of Murine CD30-L cDNA

- This example describes screening of the expression cloning library made in Example 3 with a labeled CD30/Fc fusion protein. The purified CD30/Fc fusion protein prepared in Example 1 was radioiodinated with ¹²⁵I using a commercially

available solid phase agent (ODO-GEN, Pierce). In this procedure, 5 µg of ODO-GEN were plated at the bottom of a 10 x 75 mm glass tube and incubated for twenty minutes at 4° C with 75 µl of 0.1 M sodium phosphate, pH 7.4 and 20 µl (2 mCi) Na¹²⁵I. The solution was then transferred to a second glass tube containing 5 µg of CD30/Fc in 45 µl PBS and this reaction mixture was incubated for twenty minutes at 4° C. The reaction mixture was fractionated by gel filtration on a 2 ml bed volume of Sephadex® G-25 (Sigma), and then equilibrated in RPMI 1640 medium containing 2.5% (v/v) bovine serum albumin (BSA), 0.2% (v/v) sodium azide and 20 mM Hepes, pH 7.4 binding medium. The final pool of ¹²⁵I CD30/Fc was diluted to a working stock solution of 1 x 10⁻⁷ M in binding medium, which may be stored for up to one month at 4° C without detectable loss of receptor binding activity.

Monolayers of transfected COS-7 cells made in Example 3 were assayed by slide autoradiography for expression of CD30-L using the radioiodinated CD30/Fc fusion protein. The slide autoradiographic technique was essentially as described by Gearing et al., *EMBO J.* 8:3667, 1989. Briefly, transfected COS-7 cells were washed once with binding medium (RPMI 1640 containing 25 mg/ml bovine serum albumin (BSA), 2 mg/ml sodium azide, 20 mM Hepes pH 7.2, and 50 mg/ml nonfat dry milk) and incubated for 2 hours at 4° C in binding medium containing 1 x 10⁻⁹ M ¹²⁵I-CD30/Fc fusion protein. After incubation, cells in the chambered slides were washed three times with binding buffer, followed by two washes with PBS, (pH 7.3) to remove unbound radiolabeled fusion protein.

The cells were fixed by incubating in 10% glutaraldehyde in PBS (30 minutes at room temperature), washed twice in PBS and air-dried. The slides were dipped in Kodak GTNB-2 photographic emulsion (5x dilution in water) and exposed in the dark for two to four days at room temperature in a light-proof box. The slides were developed in Kodak D19 developer, rinsed in water and fixed in Agfa G433C fixer. The slides were individually examined under a microscope at 25-40x magnification. Positive slides showing cells expressing CD30-L were identified by the presence of autoradiographic silver grains against a light background.

Eight pools, each containing approximately 2000 individual clones, were identified as positive for binding the CD30/Fc fusion protein. Two pools were titred and plated to provide plates containing approximately 200 colonies each. A replica of each breakdown pool was made and the cells were scraped to provide pooled plasmid DNA for transfection into COS-7 cells. The smaller pools were screened by slide autoradiography as described previously. Several of the breakdown pools contained clones that were positive for CD30-L as indicated by the presence of an expressed gene product capable of binding to the CD30/Fc fusion protein.

Individual colonies from two of the breakdown pools were picked from the replicas and inoculated into culture medium in individual wells of 96-well plates. Cultures were mixed by pooling rows and columns and the mixed cultures were used to prepare DNA for a final round of transfection and screening. An intersection of a positive row and a positive column identified the positive colony. DNA from the pure clone was isolated, retransfected and rescreened.

The recombinant plasmid containing murine CD30-L cDNA was recovered from the pure clone (COS-7 host cells) and transformed into *E. coli* strain DH5 α . The mammalian expression vector pDC202 containing murine CD30-L cDNA (designated pDC202-mCD30-L) was deposited in *E. coli* strain DH5 α host cells with the American Type Culture Collection, Rockville, MD (ATCC) on May 28, 1992, under accession number ATCC 69004. The deposit was made under the terms of the Budapest Treaty.

A DNA sequence for the coding region of the cDNA insert of clone pDC202-mCD30-L is presented in SEQ ID NO:18, and, the encoded amino acid sequence is presented in SEQ ID NO:19. The protein comprises an N-terminal cytoplasmic domain (amino acids 1-27), a transmembrane region (amino acids 28-48), and an extracellular, i.e., receptor-binding domain (amino acids 49-220). This protein lacks a signal peptide.

Six amino acid triplets constituting N-linked glycosylation sites are found at amino acids 56-58, 67-69, 95-97, 139-141, 175-177, and 187-189 of SEQ ID NO:19. The protein comprises no KEX2 protease processing sites.

In this particular vector construction, an ATG codon located in the Bgl II adaptors (see Example 3) is in the same reading frame as the CD30-L cDNA insert. Thus, a percentage of the transcripts may comprise the following DNA sequence upstream of the sequence of SEQ ID NO:18. The encoded amino acids are also shown, and would be fused to the N-terminus of the SEQ ID NO:19 sequence, but are not CD30-L-specific amino acids.

ATG GGC TGT GGG GCT CCT TCC CCT GAC CCA GCC (SEQ ID NO:13)

Met Gly Cys Gly Ala Pro Ser Pro Asp Pro Ala (SEQ ID NO:14)

EXAMPLE 5: Monoclonal Antibodies Directed Against CD30-L

This example illustrates the preparation of monoclonal antibodies to CD30-L. CD30-L is expressed in mammalian host cells such as COS-7 or CV1-EBNA cells and purified using CD30/Fc affinity chromatography as described herein. Purified CD30-L can be used to generate monoclonal antibodies against CD30-L using conventional techniques, for example, those techniques described in U.S. Patent 4,411,993. The immunogen may comprise a protein (or fragment thereof, such as the extracellular

domain) fused to the peptide Asp-Tyr-Lys-Asp-Asp-Asp-Lys (DYKDDDDK) (SEQ ID NO:15) (Hopp et al., *BioTechnology* 6:1204, 1988 and U.S. Patent No. 5,011,912) or fused to the Fc portion of an antibody, as described above.

- 5 Briefly, mice are immunized with CD30-L as an immunogen emulsified in complete Freund's adjuvant, and injected in amounts ranging from 10-100 µg subcutaneously or intraperitoneally. Ten to twelve days later, the immunized animals are boosted with additional CD30-L emulsified in incomplete Freund's adjuvant. Mice are periodically boosted thereafter on a weekly to bi-weekly immunization schedule. Serum samples are periodically taken by retro-orbital bleeding or tail-tip excision for testing by dot blot assay or ELISA (Enzyme-Linked Immunosorbent Assay), for
10 CD30-L antibodies.

- Following detection of an appropriate antibody titer, positive animals are provided one last intravenous injection of CD30-L in saline. Three to four days later, the animals are sacrificed, spleen cells harvested, and spleen cells are fused to a murine
15 myeloma cell line (e.g., NS1 or Ag 8.653). The latter myeloma cell line is available from the American Type Culture Collection as P3x63Ag8.653 (ATCC CRL 1580). Fusions generate hybridoma cells, which are plated in multiple microtiter plates in a HAT (hypoxanthine, aminopterin and thymidine) selective medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

- 20 The hybridoma cells are screened by ELISA for reactivity against purified CD30-L by adaptations of the techniques disclosed in Engvall et al., *Immunochem.* 8:871, 1971 and in U.S. Patent 4,703,004. A preferred screening technique is the antibody capture technique described in Beckmann et al., (*J. Immunol.* 144:4212, 1990). Positive hybridoma cells can be injected intraperitoneally into syngeneic
25 BALB/c mice to produce ascites containing high concentrations of anti-CD30-L monoclonal antibodies. Alternatively, hybridoma cells can be grown *in vitro* in flasks or roller bottles by various techniques. Monoclonal antibodies produced in mouse ascites can be purified by ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of
30 antibody to protein A or protein G can also be used, as can affinity chromatography based upon binding to CD30-L.

EXAMPLE 6: Isolation of Human CD30-L cDNA

- This example illustrates a cross-species hybridization technique which was used
35 to isolate a human CD30-L cDNA using a probe derived from the sequence of murine CD30-L. A murine CD30-L probe was produced by excising the entire cDNA insert from murine clone pDC202-mCD30-L (ATCC 69004, described in Example 4) by *Bgl*

II digestion, and ^{32}P -labeling the fragment using random primers (Boehringer-Mannheim).

5 A human peripheral blood lymphocyte (PBL) cDNA library was constructed in a phage vector ($\lambda\text{gt}10$). The PBL cells were obtained from normal human volunteers and treated with 10 ng/ml of OKT3 (an anti-CD3 antibody), and 10 ng/ml of human IL-2 (Immunex, Seattle, WA) for six days. The PBL cells were washed and stimulated with 500 ng/ml ionomycin (Calbiochem) and 10 ng/ml PMA (Sigma) for four hours. Messenger RNA was isolated from the stimulated PBL cells. cDNA synthesized on the mRNA template was packaged into $\lambda\text{gt}10$ phage vectors (Gigapak[®], Stratagene, San 10 Diego, CA) according to manufacturer's instructions. Recombinant phage were then plated on *E. coli* strain KW251 and screened using standard plaque hybridization techniques.

The murine probe was hybridized to phage cDNA in the following hybridization buffer at 37°C overnight:

15 50% Formamide
20 mM Pipes (pH 6.4)
0.8 M NaCl
2 mM EDTA
20 0.5% SDS
0.1 mg/ml salmon sperm DNA

Hybridization was followed by washing with 2X SSC, 0.1% SDS at 50°C. Positive (hybridizing) plaques were visualized by autoradiography.

25 Six of the positive plaques were purified and the inserts were isolated by PCR amplification using oligonucleotides that flank the cloning site. A partial amino acid sequence for human CD30-L was derived by determining the nucleotide sequence of a portion of one of these inserts (clone #9, about 2.0 kb in length). This partial amino acid sequence is presented in SEQ ID NO:20. The transmembrane region comprises 30 amino acids 27-48 of SEQ ID NO:20. The amino acid represented by Xaa at position 6 is most likely a methionine residue encoded by an initiation codon. This partial human sequence exhibits significant homology to an N-terminal fragment of murine CD30-L, a preliminary amino acid sequence for which is presented as SEQ ID NO:21.

35 The DNA sequence of the entire coding region of the human CD30-L clone was determined and is presented in SEQ ID NO:22 and the encoded amino acid sequence is shown in SEQ ID NO:23. The N-terminal cytoplasmic domain (amino acids 1 to 21) is followed by a transmembrane region (amino acids 22 to 43) which is followed by the

extracellular, i.e., receptor-binding domain (amino acids 44-215). This protein lacks a signal peptide. Where the partial human CD30-L of SEQ ID NO:20 differs from the full length human CD30-L amino acid sequence presented in SEQ ID NO:23, the SEQ ID NO:23 sequence is considered to be accurate. Comparison of the murine SEQ ID NO:19 and human (SEQ ID NO:23) CD30-L amino acid sequences using the above-described GAP computer program reveals 73% identity and 83% similarity between the two sequences.

Amino acid triplets that constitute potential N-linked glycosylation sites are found at positions 62-64, 90-92, 134-136, 170-172, and 182-184 in SEQ ID NO:23.

A KEX2 protease processing site is found at amino acids 72-73. If desired, these N-glycosylation processing sites may be inactivated to preclude glycosylation as described above. The KEX2 sites may be inactivated to reduce proteolysis when the CD30-L protein is expressed in yeast cells, as described above.

The products of the above-described PCR reaction (by which the cDNA insert of the positive clone was amplified) were digested with *EcoRI* and ligated into an *EcoRI*-digested vector designated pGEMBL. Plasmid pGEMBL is a derivative of the standard cloning vector pBR322 and contains a polylinker having a unique *EcoRI* site along with several other unique restriction sites. The plasmid also comprises an ampicillin resistance gene. An exemplary vector of this type is described by Dente et al., (*Nucl. Acids Res.* 11:1645, 1983).

E. coli strain DH5 α was transformed with the ligation mixture and transformants containing the desired recombinant plasmid were identified. Samples of *E. coli* DH5 α containing plasmid hCD30-L/pGEMBL were deposited with the American Type Culture Collection, Rockville, MD (ATCC) on June 24, 1992, under accession number ATCC 69020. The deposit was made under the terms of the Budapest Treaty. The deposited recombinant plasmid contains human CD30-L DNA that includes the complete coding region shown in SEQ ID NO:22.

EXAMPLE 7: Isolation of Murine and Human CD30-L DNA Encoding Additional N-Terminal Amino Acids

Because the CD30-L clones isolated in examples 4 and 6 had relatively short 5' noncoding regions and lacked stop codons upstream of the first initiation codon, isolation of CD30-L DNA comprising additional 5' sequences was attempted. An anchored PCR technique was employed, generally as described by Loh et al., *Science* 243:217 (1989) and Carrier et al., *Gene* 116:173 (1992), both of which are hereby incorporated by reference. The same procedures were employed for isolating murine and human clones, except as noted.

First strand cDNA was synthesized using a Superscript® cDNA kit (GIBCO/BRL, Gaithersburg, MD) on the following mRNA templates:

murine: 5µg total RNA from 7B9 cell line described in Example 3.

human: 2µg poly A⁺ RNA from human peripheral blood T-cells (the stimulated PBLs described in Example 6)

The primers employed in the cDNA synthesis (referred to as primers #1 hereinafter) were:

murine: 5' AGATGCTTTGACACTTG 3' (SEQ ID NO:16)

human: 5' ATCACCAGATTCCCATC 3' (SEQ ID NO:17)

Murine primer #1 is complementary to nucleotides 265-281 of SEQ ID NO:18.

Human primer #1 is complementary to nucleotides 325-341 of SEQ ID NO:22.

The reaction mixture was treated with RNAse H, then purified over a Sephadex G50 spin column (Sigma). After drying, the cDNA was resuspended in: 10 µl H₂O, 4 µl 5X terminal deoxynucleotidyl transferase (TdT) buffer (as specified by

GIBCO/BRL, Gaithersburg, MD), 4 µl 1mM dATP, and 1 µl TdT (15 units/µl). This reaction mixture was incubated at 37°C for 10 minutes to add a poly-A tail to the 3' end of the cDNA. The reaction was stopped by heating at 68°C for 15 minutes, and the mixture was applied to a Sephadex G50 spin column. The eluate was diluted to 250 µl with 10 mM Tris (pH 7.5), 1 mM EDTA. A first PCR reaction mixture was prepared by combining the first strand cDNA (tailed with adenines) with three primers in a conventional PCR reaction mixture. The primers were a first anchoring primer, a second anchoring primer, and a primer #2 (antisense).

The following reaction conditions (temperature cycles) were employed for this first PCR, and each of the PCRs described below:

94°C - 5 minutes - 1X

94°C - 0.5 minutes |

55°C - 1.5 minutes | - 30X

72°C - 2.5 minutes |

72°C - 5 minutes - 1X

The first anchoring primer contains a poly T segment that will anneal to the poly A tail added to the cDNA. This primer also inserts a NotI restriction site into the amplified DNA. The second anchoring primer, which lacks the poly T segment but is otherwise identical to the first anchoring primer, anneals (in later cycles of the reaction) to the NotI site-containing sequence inserted into the amplified DNA via the first anchoring primer.

The murine primer #2 is complementary to nucleotides 206-222 of SEQ ID NO:18. The human primer #2 is complementary to nucleotides 108-124 of SEQ ID NO:22.

5 A second PCR reaction mixture contained the products of the first PCR reaction, the 2nd anchoring primer, and primer #2. A third PCR reaction mixture contained the products of the second PCR reaction, the 2nd anchoring primer, and primer #3. The murine primer #3 contains a segment complementary to nucleotides 49-66 of SEQ ID NO:18. Human primer #3 contains a segment complementary to nucleotides 80-94 of SEQ ID NO:22. Each primer #3 also contains a segment that
10 introduces a Sall restriction site into the amplified DNA.

PCR reaction products (from PCR reaction no. 2 for human and no. 3 for murine) were separated by electrophoresis on a 1% NuSieve agarose gel (FMC Bioproducts, Rockland, ME). A PCR band comprising DNA of about 300 bp was isolated for both murine and human. The CD30-L DNA was further amplified in
15 another PCR reaction. The reaction mixture comprised:

	5 μ l	band from gel (melted at 68°C)
	10 μ l	10x buffer
	2 μ l	2nd anchoring primer
	2 μ l	primer #3
20	1 μ l	Taq DNA polymerase
	0.8 μ l	25 mM dNTP's
	79.2 μ l	dH ₂ O
	100.0 μ l	TOTAL

25 The nucleotide sequence of the reaction products was determined. The reaction products may be sequenced directly or subcloned by digesting with NotI/Sall prior to sequencing. Sequencing revealed additional DNA at the 5' end, compared to the clones of examples 4 and 6, including DNA encoding an additional 19 N-terminal amino acids for both murine and human CD30-L. DNA and encoded amino acid sequences for the
30 coding region of CD30-L DNA comprising this additional 5' coding sequence are shown in SEQ ID NO:5 and SEQ ID NO:6 (murine) and SEQ ID NO:7 and SEQ ID NO:8 (human). The additional N-terminal amino acids comprise no N-glycosylation or KEX2 protease processing sites.

The murine and human CD30-L DNAs isolated in this example were expressed
35 in CV1-EBNA cells. The molecular weight of the expressed protein, analyzed by non-reducing SDS-PAGE, was about 26,519 daltons for murine and 26,017 daltons for human CD30-L.

Although the murine and human CD30-L proteins encoded by the clones of examples 4 and 6, respectively, are truncated at the N-terminus, the encoded proteins

are biologically active in that they bind to CD30. Thus, CD30-L proteins lacking from one to all of the first 19 N-terminal amino acids shown in SEQ ID NOS:6 or 8 are biologically active CD30-L proteins of the present invention. Deletion of the first 19 amino acids of SEQ ID NOS:6 and 8 yields an amino acid sequence identical to that presented in SEQ ID NOS:19 and 23, respectively.

EXAMPLE 8: Analysis of Biological Activities of CD30-L

Cells on which CD30 expression has been previously observed were screened for a response to the recombinant CD30 ligand. Response to monoclonal antibodies that bind CD30 was also analyzed. The human cell types screened included activated T cells, three Hodgkin's lymphoma lines resembling H-RS cells with primitive B or T cell-like phenotypes, and a non-Hodgkin's lymphoma line of the large cell anaplastic lymphoma (LCAL) type.

Peripheral blood T-lymphocyte (PBT) cells were isolated by centrifugation over Histopaque (Sigma Chemical Co., St. Louis, MO) and rosetting with 2-amino-ethylisothiuronium bromide (AET)-treated sheep erythrocytes as described (Armitage et al., *Int. Immunol.* 2:1039 (1990)). The purified PBT were then cultured for 5 days in the presence of immobilized CD3 antibody and a titration of fixed CV1/EBNA cells expressing full length (membrane-bound) recombinant human CD30 ligand. In contrast to control cells transfected with vector alone, cells expressing CD30-L induced proliferation of the stimulated T-cells in a dose-dependent manner, with a maximal response observed with 2.5×10^4 CV1/EBNA cells/well. This enhanced proliferation (and other activities described below) could be blocked by the inclusion of 10 μ g/ml of soluble CD30/Fc. Proliferation of CD3-activated T cells was also seen in the presence of immobilized anti-CD30 monoclonal antibody M44, suggesting the bivalent antibody mimics ligand-induced receptor cross linking. The M44 monoclonal antibody is a mouse IgG1 generated with purified CD30-Fc as immunogen, as described further in example 12. No response to CD30-L was seen in the absence of CD3 co-stimulation.

The biological activity of CD30-L on human lymphoma cell lines known to express CD30 was investigated. The CD30⁺ human lymphoma lines tested included HDLM-2, KM-H2, L-428, and Karpas 299 cells. Culture conditions for these four cell lines are published (Drexler et al., *Leuk. Res.* 10:487 (1986); Gruss et al., *Cancer Res.* 52:3353 (1992)).

The HD-derived cell line HDLM-2 was established from a malignant pleural effusion of a 74-year-old male with endstage IVB HD (Drexler et al., 1986, *supra*; Gruss et al., 1992, *supra*). HDLM-2 is phenotypically T-cell-like (Gruss et al., 1992, *supra*). KM-H2 and L-428 are B cell-like, HD-derived lymphoma lines. The human

Karpas 299 cell line was established from blast cells in the peripheral blood of a 25-year-old white male with the diagnosis of a large cell anaplastic lymphoma (Ki-1 positive high-grade human lymphoma). The peripheral blast cells with pleomorphic nuclei resembled primitive histiocytes, which bear the surface markers CD4, CD5, HLA-DR and CD30. The Karpas 299 cell line possesses the same cytochemical, immunologic, and chromosomal profile with a 2;5 translocation as the original peripheral blood blast cells of the patient (Fischer et al., *Blood* 72:234 (1988)).

The addition of CV1/EBNA cells (10,000 cells/well) expressing recombinant human CD30-L to the HD-derived cell line HDLM-2 (50,000 cells/well) resulted in enhanced proliferation, whereas addition of control CV1/EBNA cells transfected with vector alone had minimal effect. The CD30-L-induced stimulation of HDLM-2 cell proliferation was time-dependent, with a maximal 3-4-fold enhancement observed at 72 hours. Similar results were obtained using immobilized M44 antibody, and the effect was dose-dependent. Cells cultured with an isotype-matched control monoclonal antibody showed no response. Maximal enhancement of proliferation, a five-fold increase over control cultures, was detected after stimulation with 10 µg/ml of M44 for 72 hours. Here again, the M44 CD30 monoclonal antibody has agonist characteristics and mimics properties of the ligand. In contrast to the above results, no effects of CD30-L on proliferation or viability of the KM-H2 or L-428 cells were detected, even though both lines were confirmed to be CD30⁺ by flow cytometry with M44.

A clear and dramatically different response to CD30-L was seen with the CD30⁺ non-Hodgkin lymphoma (LCAL) line Karpas 299. The addition of either CV1/EBNA cells expressing the CD30-L or M44 antibody to Karpas 299 cells (5 x 10³ cells/well) decreased the proliferation eight-fold. This effect was further analyzed with cytotoxic assays measuring ⁵¹Cr-release. Both CV1/EBNA cells expressing CD30-L and M44 antibody induced specific ⁵¹Cr release from these cells in a time and dose-dependent manner. At 18 hours, the specific release in response to CD30-L or M44 was 29.4% and 30.8%, respectively. The addition of CV1/EBNA cells transfected with vector alone, or of an isotype-matched control antibody, had no effect. Thus, in contrast to the enhanced proliferative response of the Hodgkin's lymphoma-derived HDLM-2, the response of the Karpas 299 non-Hodgkin's lymphoma line to CD30-L is cell death.

EXAMPLE 9: Northern Analysis of Murine and Human CD30-L Transcripts

Various types of cells were analyzed by Northern blotting to detect CD30-L transcripts (mRNA).

Human cells

Human PBT cells, induced with a calcium ionophore, uninduced tonsillar T cells and LPS-induced monocytes all expressed a single hybridizing transcript migrating between 18 and 28 S ribosomal RNA. IL-7-treated PBT cells, PMA treated tonsillar B cells, uninduced Jurkat or LPS activated THP-1 macrophage, and GM-CSF treated monocytes did not express CD30-L. IL-1 β induced low levels of CD30-L in monocytes. In addition, placental tissue, the promyelocytic HL60 line and two Burkitt's lymphoma B cell lines (Daudi and Raji) were also negative for expression of CD30-L transcripts. The HD-derived cell lines HDLM-2, KM-H2, and L-428, described in example 8, did not express CD30-L mRNA constitutively, or after stimulation with TPA for 24 to 72 hours or with 100 ng/mL IL-2 and TNF- α for 48 hours. Thus human CD30-L expression was detected on specifically induced T cells and monocytes/macrophages.

Murine cells

The results on human cells are mirrored in the murine system. LPS stimulated bone marrow-derived macrophage, Con A activated 7F9 T cells (similar to the 7B9 murine helper T-cell line described in examples 2 and 3) and an LPS stimulated subclone of the murine thymoma EL4 (EL4 6.1) all express a single CD30-L transcript. Unstimulated EL4 6.1 and 7F9 cells, a bone marrow-derived stromal line D11 and a thymic stromal line F4, do not express CD30-L.

EXAMPLE 10: Characterization of Recombinant CD30-L

Biochemical characteristics of the recombinant, full-length cell surface forms of murine and human CD30-L were assessed by surface radioiodinating cells transiently expressing the recombinant ligands, then immunoprecipitating the ligands with CD30/Fc (and protein G) from lysates of detergent solubilized cells. Iodoacetamide (20mM) was included in lysing and immunoprecipitation buffers to inhibit potential disulfide interchange. Washed precipitates were then displayed by SDS-PAGE with phosphorimaging. Cells transfected with vector only, or cells expressing recombinant ligand but immunoprecipitated with an isotype matched control (huIgG1), showed no bands. Under reducing conditions, the dominant product for both human and murine recombinant CD30-L is a diffuse 40 kd band. As the CD30-L protein molecular weight is 26,000 Kd, extensive use of the multiple N-linked glycosylation sites in the extracellular domains seems clear. Disulfide-linked dimers of human CD30-L appear under non-reducing conditions, and even higher oligomers, apparently disulfide-linked, are seen with murine CD30-L. Most, but not all of these are converted to monomers

upon reduction. The fact that not all oligomers were converted to monomers may reflect either differential glycosylation and/or inefficient reduction.

EXAMPLE 11: Production of a Soluble Human CD30-L Fusion Protein

5 A soluble fusion protein comprising an antibody Fc region polypeptide joined through a peptide linker to the N-terminus of a fragment of the human CD30-L extracellular domain was produced and tested for biological activity as follows. DNA encoding a soluble human CD30-L polypeptide comprising amino acids 47 (Asp) to 215 (Asp) of SEQ ID NO:23 was isolated and amplified by PCR. The PCR was
10 conducted by conventional procedures, using as the 5' primer an oligonucleotide comprising nucleotides 139-153 of SEQ ID NO:22 and a sequence containing a recognition site for BspE1. The 3' primer spanned the termination codon of CD30-L and contained the recognition sequence for Not I.

The PCR products were digested with Bsp E1 and Not I and the desired
15 fragment was ligated into an expression vector designated pDC408, which is a derivative of the pDC406 vector described above. pDC408 had been modified to contain DNA encoding (in order) 5'- murine IL-7 leader sequence - FLAG® - human IgG1 Fc domain-peptide linker.

The murine IL-7 leader sequence is described in U.S. Patent 4,965,195 and the
20 FLAG® octapeptide is described above. The Fc polypeptide is described in example 1. A peptide linker of the sequence Gly4SerGly5Ser was employed, and the soluble CD30-L encoding DNA was inserted immediately downstream of the peptide linker, in the same reading frame. 293 cells (ATCC CRL 1573; a transformed primary human embryonal kidney cell line) were transfected with the recombinant expression vector
25 and cultured to permit expression and secretion of the fusion protein. The expressed protein was purified on a protein A column.

The activity of the expressed protein was measured using an inhibition assay in which the binding of ¹²⁵I-labeled CD30/Fc protein to CD30-L expressed on the surface of transformed CV1/EBNA cells was measured. The soluble CD30-L-
30 containing fusion protein was shown to be capable of inhibiting this binding, thus indicating its ability to bind to CD30/Fc. The measured affinity of the soluble ligand for CD30/Fc was roughly equivalent to that of CD30/Fc for the cell-bound ligand.

Alternatively, an expression vector is constructed that encodes a murine IL-7 leader sequence - FLAG®- soluble CD30-L fusion protein. The Fc polypeptide and
35 peptide linker-encoding DNAs are omitted from this vector. Omitting the Fc polypeptide is advantageous in that aggregate formation is reduced. Dimers of CD30-L proteins without Fc moieties have been detected, as described in example 10. Including

an Fc polypeptide may promote formation of undesirable aggregates of oligomers of CD30-L/Fc proteins.

EXAMPLE 12: Antibodies That Bind CD30

- 5 To generate monoclonal antibodies against the human CD30 antigen, CB6F1 mice (purchased from Jackson Laboratories, Bar Harbor, ME) were boosted twice intradermally with 10 μ g CD30/Fc in Ribi adjuvant (Ribi Immunochem Research, Hamilton, MI). The soluble human CD30/Fc fusion protein employed as the immunogen was produced as described in example 1. One week after the second
- 10 boost, peroxidase dot blot assays using CD30/Fc showed a significant ($>1/100$) titer of anti-CD30 antibody in the serum. One week later, animals were boosted intravenously (IV) with 3 μ g CD30/Fc into the tail vein. Three days later, spleen was removed and spleen cells were fused to the X63-Ag8.653 mouse myeloma cell line (Kearney et al., *J. Immunol.* 123:1548, 1979) by standard methods using a 50% polyethylene
- 15 glycol/dimethyl sulfoxide solution (Sigma). Hybridoma cultures were established in 96-well plates (Costar, Cambridge, MA). Ten days later, culture supernatants were screened by an antigen capture assay using 125 I-CD30/Fc. Ninety-six-well plates were coated overnight with goat-antimouse serum (Zymed, San Francisco, CA) and blocked with 3% bovine serum albumin (BSA; Sigma); 50 μ L of culture supernatant was
- 20 incubated for 1 hour at room temperature. After three washes with phosphate-buffered saline (PBS), plates were incubated with 125 I-CD30/Fc for 1 hour and then washed with PBS again before being placed on film for overnight exposure. Positive wells were checked for reactivity with huIgG by performing an anti-biotin complex assay. Hybridoma cell lines reactive with HuIgG:horseradish peroxidase-CD30 were cloned.
- 25 Positive supernatants were also tested by flow cytometry using CD30-expressing cells or CD30-transfected CV-1/EBNA cells.

- Two human anti-CD30 monoclonal antibodies designated M44 and M67 (mouse IgG₁ isotype) were purified from spent bulk culture supernatants from two hybridoma cell lines produced above and grown in roller bottles. Antibodies were
- 30 purified on a protein A affinity matrix using an automated purification system (BioRad MAPS system, Hercules, CA). Antibody concentration was determined by absorbance at 280 nm and purity assessed by sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis and silver staining. Antibody concentrations were adjusted to 1 mg/mL and aliquots of purified antibody were stored frozen at -20°C in 0.05 mol/L citrate
- 35 buffer (pH 7.0).

EXAMPLE 13: Analysis of Biological Activities of CD30-L

Further studies of the response of cells expressing CD30 to the recombinant CD30 ligand were conducted. Response to monoclonal antibodies reactive with CD30 was also analyzed. These studies are similar to those described in example 8, but

5 include additional cell lines and antibodies. The human cell types screened included activated T cells, four Hodgkin's lymphoma lines resembling H-RS cells with primitive B or T cell-like phenotypes, non-Hodgkin's lymphoma lines of the large cell anaplastic lymphoma (LCAL) type, and a T-cell leukemia (T-ALL) line.

10 Activated T-Cells.

Peripheral blood T-lymphocyte (PBT) cells were isolated from normal healthy human donors by centrifugation over Histopaque (Sigma Chemical Co., St. Louis, MO) and rosetting with 2-aminoethylisothiuronium bromide (AET)-treated sheep erythrocytes as described (Armitage et al., *Int. Immunol.* 2:1039 (1990)).

- 15 Contaminating monocytes were removed by plastic adherence for 1 hour at 37°C. The resulting T-cell preparations were greater than 98% CD3⁺, as determined by flow cytometric analysis. For activation of T-cells to induce CD30 expression, 96-well plates were coated with 10 µg/mL OKT3 (an anti-CD3-antibody; ATCC-CRL8001) in 50 mmol/L Tris buffer (pH 8.5) and washed twice with PBS. The purified T-cells (1 x
- 20 10⁵ cells/well) were then cultured for 72 hours in the presence of the immobilized anti-CD3 antibody and one of the following: a titration of fixed CV1/EBNA cells expressing full length (membrane-bound) recombinant human CD30 ligand; a titration of CV1/EBNA cells transformed with the empty expression vector alone; medium alone; a titration of immobilized anti-CD30 monoclonal antibody M44 or M67
- 25 (described in example 12); or a titration of an immobilized isotype-matched control antibody. The CV1/EBNA cells employed in the assay were transfected using the diethyl aminoethyl (DEAE) /Dextran method with either vector alone or a CD30L cDNA containing expression vector (CV-1/CD30L), and then fixed at 2 days posttransfection with 1% paraformaldehyde for 5 minutes at 25°C. The transformed CV1/EBNA cells
- 30 were employed in the assay at 5 x 10⁴ cells/well; the antibodies at a concentration of 10 µg/ml.

Cultures were pulsed with 1 µCi/well ³H-thymidine (³H.TdR; 25 Ci/mmol: Amersham, Arlington Heights, IL) for the final 12 hours of culture. Cells were harvested and incorporated cpm determined by tritium-sensitive avalanche gas ionization

35 detection on a Matrix 96 Beta Counter (Packard, Meriden, CT).

Cells expressing murine or human CD30-L induced proliferation of the stimulated T-cells whereas no proliferative response was induced by medium or

CV1/EBNA cells transformed with the empty vector. Proliferation of CD3-activated T-cells was also induced by the immobilized anti-CD30 monoclonal antibodies M44 and M67, suggesting the bivalent antibody mimics ligand-induced receptor cross linking. The activated T-cells did not respond to the irrelevant control antibody.

5

Lymphoma Cell Lines

The biological activity of CD30-L and anti-CD30 antibodies on human lymphoma cell lines known to express CD30 was investigated. The CD30⁺ EBV-human lymphoma lines tested included the Hodgkins Disease (HD) derived lines HDLM-2, KM-H2, L-428, and L-540, and several LCAL-type non-Hodgkins' lymphoma lines. These cell lines and appropriate culture conditions are described in Drexler et al., *Leuk. Res.* 10:487 (1986); Gruss et al., *Cancer Res.* 52:3353 (1992); Kamesaki et al., *Blood* 68:285 (1986); Schaadt et al., *Int. J. Cancer* 26:723 (1980); and Diehl et al., *J. Cancer Res. Clin. Oncol.* 101:111 (1981).

15 The HD-derived cell line HDLM-2 was established from a malignant pleural effusion of a 74-year-old male with endstage IVB nodular sclerosis (NS) HD (Drexler et al., 1986, *supra*; Gruss et al., 1992, *supra*) and is phenotypically T-cell-like (Gruss et al., 1992, *supra*). KM-H2 and L-428 are B cell-like, HD-derived lymphoma lines. The L-428 cell line was derived from a malignant pleural effusion of a 37-year-old woman with endstage IVB NS HD; the KM-H2 cell line from a malignant pleural effusion of a 37-year-old man with stage IV mixed cellularity (MC) HD. The L-540 cell line, which is phenotypically T-cell-like, was derived from the bone marrow of a 20-year-old woman with stage IVB NS HD.

The human large cell anaplastic lymphoma cell line Karpas 299 was established from blast cells in the peripheral blood of a 25-year-old white male with the diagnosis of CD30⁺ high-grade LCAL. The peripheral blast cells with pleomorphic nuclei resembled primitive histiocytes, which bear the surface markers CD4, CD5, epithelial membrane antigen (EMA), HLA-DR and CD30. The Karpas 299 cell line possesses the same cytochemical, immunologic, morphologic and chromosomal profile with a 2;5 translocation as the original peripheral blood blast cells of the patient (Fischer et al., *Blood* 72:234 (1988)). Seven additional permanent LCAL cell lines employed in the study were established from primary CD30⁺ LCAL tumors, and resemble the malignant lymphoma clone of the primary LCAL patients.

Proliferative responses of the cell lines to CD30-L and anti-CD30 MAbs was analyzed in a thymidine incorporation assay similar to that described in example 8, as follows. A total of 5×10^4 HDLM-2, L-540, L-428, KM-H2 or LCAL (e.g., Karpas 299) cells were cultured for 72 hours with 1×10^4 CV-1/EBNA cells transfected with

empty vector, with human or murine CD30-L encoding vectors, or with 10 µg/mL immobilized anti-CD30 MAbs M44, M67, and Ki-1 or isotype control MAb. The Ki-1 MAb (included in all but the assay on L_{CAL} cells) was purchased from Dako Corporation, Santa Barbara, CA. Tritiated-thymidine incorporation was determined after 72 hours.

No effects of CD30-L, M44, M67, or Ki-1 on proliferation or viability of the "B-cell-like" KM-H2 or L-428 cells were detected, even though both lines were confirmed to be CD30⁺ by flow cytometry with anti-CD30 MAbs. In contrast, proliferation of the "T-cell-like" HD-derived cell lines HDLM-2 and L-540 was enhanced after addition of CD30-L or anti-CD30 MoAbs M44 and M67. Both murine and human CD30-L, expressed on CV-1 EBNA cells, induced a twofold to fivefold enhancement of ³H-thymidine uptake by HDLM-2 and L-540 H-RS cells compared with cells cultured with medium or CV-1/EBNA cells containing only the empty vector. Also, immobilized anti-CD30 MAbs M44 and M67 enhanced proliferation of HDLM-2 and L-540 cells threefold to eightfold. In contrast, the anti-CD30 MAb Ki-1 did not induce proliferation of HDLM-2 and L-540 cells above that induced by isotype-matched control antibody.

The CD30-L- and M44/M67-induced stimulation of HDLM-2 and L-540 cell proliferation was time- and dose-dependent. CD30-L induced maximal proliferation after 72 hours of culture. The MAbs M44 and M67 had maximal effects at concentrations of 10 µg/mL for a culture period of 48 to 72 hours. The enhanced proliferation effect of the CD30-L or the agonistic MAbs appeared to be specific because it could be blocked by the addition of 50-fold excess of soluble CD30/Fc protein. Here again, the M44 and M67 anti-CD30 monoclonal antibodies have agonist characteristics and mimic properties of the ligand.

A different response to CD30-L, M44 and M67 was seen with the CD30⁺ non-Hodgkin lymphoma (L_{CAL}) line Karpas 299. The addition of either CV1/EBNA cells expressing the CD30-L, or the M44 or M67 antibodies to Karpas 299 cells resulted in a threefold to sixfold reduction in proliferation in comparison to cells cultured with CV-1/EBNA cells transfected with the vector alone, isotype-matched control MAb, or medium. A significant reduction of ³H-thymidine uptake by Karpas 299 cells is measurable after 24 hours in culture with CD30-L, M44, or M67. The reduction of proliferation was time-dependent (being minimal 72 hours after initiation of the cultures), dose-dependent, and could be almost completely reversed by the addition of 50-fold excess of soluble CD30/Fc. Only one of the seven additional L_{CAL} cell lines did not show an alteration of proliferation after the addition of membrane-expressed CD30-L or anti-CD30 MAbs M44 and M67. A 30% to 70% reduction of

proliferation in response to CD30-L, M44, and M67, in comparison to the controls, was seen for the other LCAL lines.

T-Cell Leukemia Cell Line

- 5 The proliferative response of an adult T-cell leukemia (T-ALL) cell line designated KE-37 to CD30-L (human only) and antibodies M44 and M67 was analyzed in the above-described thymidine incorporation assay. The recombinant CD30-L and both anti-CD30 antibodies induced enhanced proliferation of the KE-37 cells. No proliferative response was seen for any of the controls (including the antibody Ki-1).

CLAIMS

What is claimed is:

1. An isolated DNA sequence encoding a CD30-L polypeptide, wherein said CD30-L comprises an amino acid sequence selected from the group consisting of amino acids 1-220 of SEQ ID NO:19, amino acids 1-215 of SEQ ID NO:23, amino acids 1-239 of SEQ ID NO:6, and amino acids 1-234 of SEQ ID NO:8.
2. An isolated DNA sequence encoding a soluble CD30-L polypeptide capable of binding CD30, wherein said CD30-L comprises an amino acid sequence selected from the group consisting of amino acids 49-220 of SEQ ID NO:19 and amino acids z-215 of SEQ ID NO:23, wherein z is selected from the group consisting of 44, 45, 46, and 47.
3. An isolated DNA comprising a DNA sequence that encodes an Fc polypeptide derived from an antibody, fused directly or through a peptide linker-encoding sequence to the 5' end of a soluble CD30-L-encoding DNA according to claim 2.
4. A DNA sequence according to claim 2, wherein said DNA sequence comprises a nucleotide sequence selected from the group consisting of nucleotides 130-645, 133-645, 136-645, and 139-645 of SEQ ID NO:22.
5. An isolated DNA capable of hybridizing to a DNA sequence of claim 1 under highly stringent conditions, wherein said isolated DNA encodes a CD30-L polypeptide capable of binding CD30.
6. An isolated DNA sequence encoding a CD30-L polypeptide capable of binding CD30, wherein said CD30-L comprises an amino acid sequence selected from the group consisting of amino acids x to 239 of SEQ ID NO:6, wherein x is 1-19, and amino acids y to 234 of SEQ ID NO:8, wherein y is 1-19.
7. An expression vector comprising a DNA sequence according to claim 1.
8. An expression vector comprising a DNA sequence according to claim 2.

9. An expression vector comprising a DNA sequence according to claim 3.
10. An expression vector comprising a DNA sequence according to claim 5.
11. A process for preparing a CD30-L polypeptide, comprising culturing a host cell transformed with a vector according to claim 7 under conditions promoting expression of CD30-L, and recovering the CD30-L polypeptide.
12. A process for preparing a CD30-L polypeptide, comprising culturing a host cell transformed with a vector according to claim 8 under conditions promoting expression of CD30-L and recovering the CD30-L polypeptide.
13. A process for preparing a soluble CD30-L/Fc fusion protein, comprising culturing a host cell transformed with a vector according to claim 9 under conditions promoting expression of CD30-L/Fc, and recovering the CD30-L/Fc polypeptide.
14. A process for preparing a CD30-L polypeptide, comprising culturing a host cell transformed with a vector according to claim 10 under conditions promoting expression of CD30-L, and recovering the CD30-L polypeptide.
15. A substantially homogeneous purified biologically active CD30-L protein, wherein said CD30-L is selected from the group consisting of murine CD30-L comprising the N-terminal amino acid sequence Met-Gln-Val-Gln-Pro-Gly-Ser-Val-Ala-Ser-Pro-Trp or Met-Glu-Pro-Gly-Leu-Gln-Gln-Ala-Gly-Ser-Cys-Gly, and human CD30-L comprising the N-terminal amino acid sequence Met-His-Val-Pro-Ala-Gly-Ser-Val-Ala-Ser-His-Leu or Met-Asp-Pro-Gly-Leu-Gln-Gln-Ala-Leu-Asn-Gly-Met.
16. A purified CD30-L according to claim 15, wherein said CD30-L comprises an amino acid sequence selected from the group consisting of amino acids 1-220 of SEQ ID NO:19, amino acids 1-215 of SEQ ID NO:23, amino acids 1-239 of SEQ ID NO:6, and amino acids 1-234 of SEQ ID NO:8.
17. A substantially homogeneous soluble CD30-L polypeptide, wherein said soluble CD30-L comprises an amino acid sequence selected from the group consisting of amino acids 49-220 of SEQ ID NO:19 and amino acids z-215 of SEQ ID NO:23, wherein z is selected from the group consisting of 44, 45, 46, and 47.

18. A substantially homogeneous purified CD30-L protein capable of binding CD30, wherein said CD30-L is encoded by a DNA sequence that will hybridize to the nucleotide sequence presented in SEQ ID NO:19 or SEQ ID NO:23 under severely stringent conditions.

19. A CD30-L protein according to claim 18, wherein said protein is a soluble human CD30-L protein.

20. Purified CD30-L according to claim 18, wherein said CD30-L comprises an amino acid sequence selected from the group consisting of amino acids x to 239 of SEQ ID NO:6, wherein x is 1-19, and amino acids y to 234 of SEQ ID NO:8, wherein y is 1-19.

21. A fusion protein comprising a soluble human CD30-L according to claim 19 and an Fc polypeptide derived from an antibody.

22. An antibody immunoreactive with a CD30-L polypeptide according to claim 18.

23. An antibody according to claim 22 wherein said antibody is a monoclonal antibody.

24. An isolated nucleic acid molecule comprising a sequence of at least about 14 nucleotides of the DNA sequence of SEQ ID NO:5 or SEQ ID NO:7, or the DNA or RNA complement thereof.

25. A conjugate comprising a diagnostic or therapeutic agent attached to a CD30-L polypeptide according to claim 18.

26. A conjugate comprising a diagnostic or therapeutic agent attached to a soluble CD30-L polypeptide according to claim 19.

27. A method of delivering a diagnostic or therapeutic agent to CD30⁺ cells, comprising contacting said cells with a conjugate according to claim 25.

28. A method of delivering a diagnostic or therapeutic agent to CD30⁺ cells, comprising contacting said cells with a conjugate according to claim 26.

29. A method according to claim 28, wherein said cells are malignant and said conjugate is administered in an effective amount to a human afflicted with said malignant cells.

30. A method according to claim 29, wherein said cells are CD30⁺ lymphoma cells.

31. A method of treating large cell anaplastic lymphoma (LCAL), comprising administering an effective amount of a soluble CD30-L according to claim 19 to a human afflicted with LCAL.

ABSTRACT OF THE DISCLOSURE

There is disclosed a polypeptide (CD30-L) and DNA sequences, vectors and transformed host cells useful in providing CD30-L polypeptides. The CD30-L polypeptide binds to the receptor known as CD30, which is expressed on a number of cell types, among which are Hodgkin's Disease tumor cells, large cell anaplastic lymphoma cells, adult T-cell leukemia (T-ALL) cells, and a number of other malignant cell types. CD30-L polypeptides find use as carriers for delivering diagnostic and cytotoxic agents to cells expressing the CD30 receptor.

Immunex Corporation

2804-E

DECLARATION AND POWER OF ATTORNEY

As the below-named inventors, we declare that we are the original, first, and sole inventors of the subject matter which is claimed in the specification identified below and for which a patent is sought on the invention as titled therein. We hereby state that we have reviewed and understand the contents of said specification including the claims. We acknowledge the duty to disclose information which is known to us to be material to the patentability of this application in accordance with 37 C.F.R. 1.56.

Inventor: GOODWIN, Raymond G.
Address: 3322 8th Avenue West

Seattle, Washington 98119
U.S.A.

Residence: Seattle, Washington, U.S.A.
Citizenship: U.S.A.

Inventor: SMITH, Craig A.
Address: 2405 - 5th West

Seattle, Washington 98119
U.S.A.

Residence: Seattle, Washington, U.S.A.
Citizenship: U.S.A.

Inventor: ARMITAGE, Richard J.
Address: 5133 Eagle Harbor Drive

Bainbridge Island, Washington 98110
U.S.A.

Residence: Bainbridge Island, Washington, U.S.A.
Citizenship: United Kingdom

Inventor: GRUSS, Hans-Juergen
Address: 930 Blue Heron N.E. Avenue

Bainbridge Island, Washington 98110
U.S.A.

Residence: Bainbridge Island, Washington, U.S.A.
Citizenship: German

Title of the Invention: NOVEL CYTOKINE THAT BINDS CD30

USSN: 08/225,989, filed on April 12, 1994.

() There are no earlier-filed U. S. applications of which priority benefit is claimed.

(X) We hereby claim the benefit under 35 U.S.C. 120 of the United States applications listed below, and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States applications in the manner provided by the first

Immunex Corporation
Declaration and Power of Attorney
Docket No. 2804-E

paragraph of 35 U.S.C. 112, we acknowledge the duty to disclose material information as defined in 37 C.F.R. 1.56 which occurred between the filing date of the prior applications and the filing date of this application:

<u>USSN:</u>	<u>Filed:</u>	<u>Status:</u>
07/966,775	October 27, 1992	Pending
07/907,224	July 1, 1992	Abandoned
07/899,660	June 15, 1992	Abandoned
07/892,459	June 2, 1992	Abandoned
07/889,717	May 26, 1992	Abandoned

POWER OF ATTORNEY

The power to prosecute this application and transact all business in the Patent and Trademark Office connected herewith is hereby granted to the following attorneys and agents:

Scott G. Hallquist Registration No. 30,641	Christopher L. Wight Registration No. 31,680
Kathryn A. Seese Registration No. 32,172	Patricia Anne Perkins Registration No. 34,693
Charlene A. Launer Registration No. 33,035	Stephen L. Malaska Registration No. 32,655

Send all correspondence and direct telephone calls to:

Legal Affairs Department
Immunex Corporation
51 University Street
Seattle, WA 98101
Telephone: (206) 587-0430

We further declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of

Immunex Corporation
Declaration and Power of Attorney
Docket No. 2804-E

Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor:

Date Signed:

Raymond A. Goodwin
Raymond G. Goodwin

6/14/94

Craig A. Smith
Craig A. Smith

6/14/94

Richard J. Armitage
Richard J. Armitage

7/8/94

Hans-Jürgen Gruss
Hans-Jürgen Gruss

June 13, 1994

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Goodwin, Raymond G.
Smith, Craig A.
Armitage, Richard J.
Gruss, Hans-Jurgen
- (ii) TITLE OF INVENTION: Novel Cytokine That Binds CD30
- (iii) NUMBER OF SEQUENCES: 23
- (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Kathryn A. Seese, Immunex Corporation
(B) STREET: 51 University Street
(C) CITY: Seattle
(D) STATE: Washington
(E) COUNTRY: USA
(F) ZIP: 98101
- (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: Apple Macintosh
(C) OPERATING SYSTEM: Apple 7.1
(D) SOFTWARE: Microsoft Word, Version 5.1a
- (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER: --to be assigned--
(B) FILING DATE: 12 APRIL 1994
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: US 07/966,775
(B) FILING DATE: 27-OCT-1992
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: US 907,224
(B) FILING DATE: 01-JUL-1992
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: US 899,660
(B) FILING DATE: 15-JUN-1992
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: US 892,459
(B) FILING DATE: 02-JUN-1992
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: US 889,717
(B) FILING DATE: 26-MAY-1992
- (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Seese, Kathryn A.
(B) REGISTRATION NUMBER: 32,172
(C) REFERENCE/DOCKET NUMBER: 2804-E

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (206) 587-0430
(B) TELEFAX: (206) 233-0644
(C) TELEX: 756822

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1788 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

(B) CLONE: huCD30

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..1788

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG	CGC	GTC	CTC	CTC	GCC	GCG	CTG	GGA	CTG	CTG	TTC	CTG	GGG	GCG	CTA	48
Met	Arg	Val	Leu	Leu	Ala	Ala	Leu	Gly	Leu	Leu	Phe	Leu	Gly	Ala	Leu	
1				5				10					15			
CGA	GCC	TTC	CCA	CAG	GAT	CGA	CCC	TTC	GAG	GAC	ACC	TGT	CAT	GGA	AAC	96
Arg	Ala	Phe	Pro	Gln	Asp	Arg	Pro	Phe	Glu	Asp	Thr	Cys	His	Gly	Asn	
			20					25					30			
CCC	AGC	CAC	TAC	TAT	GAC	AAG	GCT	GTC	AGG	AGG	TGC	TGT	TAC	CGC	TGC	144
Pro	Ser	His	Tyr	Tyr	Asp	Lys	Ala	Val	Arg	Arg	Cys	Cys	Tyr	Arg	Cys	
			35				40					45				
CCC	ATG	GGG	CTG	TTC	CCG	ACA	CAG	CAG	TGC	CCA	CAG	AGG	CCT	ACT	GAC	192
Pro	Met	Gly	Leu	Phe	Pro	Thr	Gln	Gln	Cys	Pro	Gln	Arg	Pro	Thr	Asp	
	50					55					60					
TGC	AGG	AAG	CAG	TGT	GAG	CCT	GAC	TAC	TAC	CTG	GAT	GAG	GCC	GAC	CGC	240
Cys	Arg	Lys	Gln	Cys	Glu	Pro	Asp	Tyr	Tyr	Leu	Asp	Glu	Ala	Asp	Arg	
	65				70					75					80	
TGT	ACA	GCC	TGC	GTG	ACT	TGT	TCT	CGA	GAT	GAC	CTC	GTG	GAG	AAG	ACG	288
Cys	Thr	Ala	Cys	Val	Thr	Cys	Ser	Arg	Asp	Asp	Leu	Val	Glu	Lys	Thr	
			85						90					95		
CCG	TGT	GCA	TGG	AAC	TCC	TCC	CGT	GTC	TGC	GAA	TGT	CGA	CCC	GGC	ATG	336
Pro	Cys	Ala	Trp	Asn	Ser	Ser	Arg	Val	Cys	Glu	Cys	Arg	Pro	Gly	Met	
			100					105					110			

TTC	TGT	TCC	ACG	TCT	GCC	GTC	AAC	TCC	TGT	GCC	CGC	TGC	TTC	TTC	CAT	384
Phe	Cys	Ser	Thr	Ser	Ala	Val	Asn	Ser	Cys	Ala	Arg	Cys	Phe	Phe	His	
	115						120					125				
TCT	GTC	TGT	CCG	GCA	GGG	ATG	ATT	GTC	AAG	TTC	CCA	GGC	ACG	CGC	CAG	432
Ser	Val	Cys	Pro	Ala	Gly	Met	Ile	Val	Lys	Phe	Pro	Gly	Thr	Ala	Gln	
	130						135					140				
AAG	AAC	ACG	GTC	TGT	GAG	CCG	GCT	TCC	CCA	GGG	GTC	AGC	CCT	GCC	TGT	480
Lys	Asn	Thr	Val	Cys	Glu	Pro	Ala	Ser	Pro	Gly	Val	Ser	Pro	Ala	Cys	
	145				150					155					160	
GCC	AGC	CCA	GAG	AAC	TGC	AAG	GAA	CCC	TCC	AGT	GGC	ACC	ATC	CCC	CAG	528
Ala	Ser	Pro	Glu	Asn	Cys	Lys	Glu	Pro	Ser	Ser	Gly	Thr	Ile	Pro	Gln	
			165						170					175		
GCC	AAG	CCC	ACC	CCG	GTG	TCC	CCA	GCA	ACC	TCC	AGT	GCC	AGC	ACC	ATG	576
Ala	Lys	Pro	Thr	Pro	Val	Ser	Pro	Ala	Thr	Ser	Ser	Ala	Ser	Thr	Met	
			180						185					190		
CCT	GTA	AGA	GGG	GGC	ACC	CGC	CTC	GCC	CAG	GAA	GCT	GCT	TCT	AAA	CTG	624
Pro	Val	Arg	Gly	Gly	Thr	Arg	Leu	Ala	Gln	Glu	Ala	Ala	Ser	Lys	Leu	
			195				200						205			
ACG	AGG	GCT	CCC	GAC	TCT	CCC	TCC	TCT	GTG	GGA	AGG	CCT	AGT	TCA	GAT	672
Thr	Arg	Ala	Pro	Asp	Ser	Pro	Ser	Ser	Val	Gly	Arg	Pro	Ser	Ser	Asp	
	210					215					220					
CCA	GGT	CTG	TCC	CCA	ACA	CAG	CCA	TGC	CCA	GAG	GGG	TCT	GGT	GAT	TGC	720
Pro	Gly	Leu	Ser	Pro	Thr	Gln	Pro	Cys	Pro	Glu	Gly	Ser	Gly	Asp	Cys	
	225					230				235				240		
AGA	AAG	CAG	TGT	GAG	CCC	GAC	TAC	TAC	CTG	GAC	GAG	GCC	GGC	CGC	TGC	768
Arg	Lys	Gln	Cys	Glu	Pro	Asp	Tyr	Tyr	Leu	Asp	Glu	Ala	Gly	Arg	Cys	
			245						250					255		
ACA	GCC	TGC	GTG	AGC	TGT	TCT	CGA	GAT	GAC	CTT	GTG	GAG	AAG	ACG	CCA	816
Thr	Ala	Cys	Val	Ser	Cys	Ser	Arg	Asp	Asp	Leu	Val	Glu	Lys	Thr	Pro	
			260					265					270			
TGT	GCA	TGG	AAC	TCC	TCC	CGC	ACC	TGC	GAA	TGT	CGA	CCT	GGC	ATG	ATC	864
Cys	Ala	Trp	Asn	Ser	Ser	Arg	Thr	Cys	Glu	Cys	Arg	Pro	Gly	Met	Ile	
		275					280					285				
TGT	GCC	ACA	TCA	GCC	ACC	AAC	TCC	TGT	GCC	CGC	TGT	GTC	CCC	TAC	CCA	912
Cys	Ala	Thr	Ser	Ala	Thr	Asn	Ser	Cys	Ala	Arg	Cys	Val	Pro	Tyr	Pro	
		290				295					300					
ATC	TGT	GCA	GGA	GAG	ACG	GTC	ACC	AAG	CCC	CAG	GAT	ATG	GCT	GAG	AAG	960
Ile	Cys	Ala	Gly	Glu	Thr	Val	Thr	Lys	Pro	Gln	Asp	Met	Ala	Glu	Lys	
	305					310				315				320		
GAC	ACC	ACC	TTT	GAG	GCG	CCA	CCC	CTG	GGG	ACC	CAG	CCG	GAC	TGC	AAC	1008
Asp	Thr	Thr	Phe	Glu	Ala	Pro	Pro	Leu	Gly	Thr	Gln	Pro	Asp	Cys	Asn	
			325						330					335		
CCC	ACC	CCA	GAG	AAT	GGC	GAG	GCG	CCT	GCC	AGC	ACC	AGC	CCC	ACT	CAG	1056
Pro	Thr	Pro	Glu	Asn	Gly	Glu	Ala	Pro	Ala	Ser	Thr	Ser	Pro	Thr	Gln	
			340					345					350			

AGC	TTG	CTG	GTG	GAC	TCC	CAG	GCC	AGT	AAG	ACG	CTG	CCC	ATC	CCA	ACC	1104
Ser	Leu	Leu	Val	Asp	Ser	Gln	Ala	Ser	Lys	Thr	Leu	Pro	Ile	Pro	Thr	
	355						360					365				
AGC	GCT	CCC	GTC	GCT	CTC	TCC	TCC	ACG	GGG	AAG	CCC	GTT	CTG	GAT	GCA	1152
Ser	Ala	Pro	Val	Ala	Leu	Ser	Ser	Thr	Gly	Lys	Pro	Val	Leu	Asp	Ala	
	370					375					380					
GGG	CCA	GTG	CTC	TTC	TGG	GTG	ATC	CTG	GTG	TTG	GTT	GTG	GTG	GTC	GGC	1200
Gly	Pro	Val	Leu	Phe	Trp	Val	Ile	Leu	Val	Leu	Val	Val	Val	Val	Gly	
	385				390					395					400	
TCC	AGC	GCC	TTC	CTC	CTG	TGC	CAC	CGG	AGG	GCC	TGC	AGG	AAG	CGA	ATT	1248
Ser	Ser	Ala	Phe	Leu	Leu	Cys	His	Arg	Arg	Ala	Cys	Arg	Lys	Arg	Ile	
			405						410					415		
CGG	CAG	AAG	CTC	CAC	CTG	TGC	TAC	CCG	GTC	CAG	ACC	TCC	CAG	CCC	AAG	1296
Arg	Gln	Lys	Leu	His	Leu	Cys	Tyr	Pro	Val	Gln	Thr	Ser	Gln	Pro	Lys	
			420						425					430		
CTA	GAG	CTT	GTG	GAT	TCC	AGA	CCC	AGG	AGG	AGC	TCA	ACG	CAG	CTG	AGG	1344
Leu	Glu	Leu	Val	Asp	Ser	Arg	Pro	Arg	Arg	Ser	Ser	Thr	Gln	Leu	Arg	
	435						440					445				
AGT	GGT	GCG	TCG	GTG	ACA	GAA	CCC	GTC	GCG	GAA	GAG	CGA	GGG	TTA	ATG	1392
Ser	Gly	Ala	Ser	Val	Thr	Glu	Pro	Val	Ala	Glu	Ala	Arg	Gly	Leu	Met	
	450					455						460				
AGC	CAG	CCA	CTG	ATG	GAG	ACC	TGC	CAC	AGC	GTG	GGG	GCA	GCC	TAC	CTG	1440
Ser	Gln	Pro	Leu	Met	Glu	Thr	Cys	His	Ser	Val	Gly	Ala	Ala	Tyr	Leu	
	465				470					475					480	
GAG	AGC	CTG	CCG	CTG	CAG	GAT	GCC	AGC	CCG	GCC	GGG	GGC	CCC	TGG	TCC	1488
Glu	Ser	Leu	Pro	Leu	Gln	Asp	Ala	Ser	Pro	Ala	Gly	Gly	Pro	Ser	Ser	
			485						490					495		
CCC	AGG	GAC	CTT	CCT	GAG	CCC	CGG	GTG	TCC	ACG	GAG	CAC	ACC	AAT	AAC	1536
Pro	Arg	Asp	Leu	Pro	Glu	Pro	Arg	Val	Ser	Thr	Glu	His	Thr	Asn	Asn	
		500						505					510			
AAG	ATT	GAG	AAA	ATC	TAC	ATC	ATG	AAG	GCT	GAC	ACC	GTG	ATC	GTG	GGG	1584
Lys	Ile	Glu	Lys	Ile	Tyr	Ile	Met	Lys	Ala	Asp	Thr	Val	Ile	Val	Gly	
	515						520					525				
ACC	GTG	AAG	GCT	GAG	CTG	CCG	GAG	GGC	CGG	GGC	CTG	GCG	GGG	CCA	GCA	1632
Thr	Val	Lys	Ala	Glu	Leu	Pro	Glu	Gly	Arg	Gly	Leu	Ala	Gly	Pro	Ala	
	530					535					540					
GAG	CCC	GAG	TTG	GAG	GAG	GAG	CTG	GAG	GCG	GAC	CAT	ACC	CCC	CAC	TAC	1680
Glu	Pro	Glu	Leu	Glu	Glu	Glu	Leu	Glu	Ala	Asp	His	Thr	Pro	His	Tyr	
	545				550				555						560	
CCC	GAG	CAG	GAG	ACA	GAA	CCG	CCT	CTG	GGC	AGC	TGC	AGC	GAT	GTC	ATG	1728
Pro	Glu	Gln	Glu	Thr	Glu	Pro	Pro	Leu	Gly	Ser	Cys	Ser	Asp	Val	Met	
			565						570					575		
CTC	TCA	GTG	GAA	GAG	GAA	GGG	AAA	GAA	GAC	CCC	TTG	CCC	ACA	GCT	GCC	1776
Leu	Ser	Val	Glu	Glu	Glu	Gly	Lys	Glu	Asp	Pro	Leu	Pro	Thr	Ala	Ala	
		580						585					590			

TCT GGA AAG TGA
Ser Gly Lys
595

1788

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 595 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Arg	Val	Leu	Leu	Ala	Ala	Leu	Gly	Leu	Leu	Phe	Leu	Gly	Ala	Arg	1	5	10	15
Arg	Ala	Phe	Pro	Gln	Asp	Arg	Pro	Phe	Glu	Asp	Thr	Cys	His	Gly	Asn	20	25	30	
Pro	Ser	His	Tyr	Tyr	Asp	Lys	Ala	Val	Arg	Arg	Cys	Cys	Tyr	Arg	Cys	35	40	45	
Pro	Met	Gly	Leu	Phe	Pro	Thr	Gln	Gln	Cys	Pro	Gln	Arg	Pro	Thr	Asp	50	55	60	
Cys	Arg	Lys	Gln	Cys	Glu	Pro	Asp	Tyr	Tyr	Leu	Asp	Glu	Ala	Asp	Arg	65	70	75	80
Cys	Thr	Ala	Cys	Val	Thr	Cys	Ser	Arg	Asp	Asp	Leu	Val	Glu	Lys	Thr	85	90	95	
Pro	Cys	Ala	Trp	Asn	Ser	Ser	Arg	Val	Cys	Glu	Cys	Arg	Pro	Gly	Met	100	105	110	
Phe	Cys	Ser	Thr	Ser	Ala	Val	Asn	Ser	Cys	Ala	Arg	Cys	Phe	Phe	His	115	120	125	
Ser	Val	Cys	Pro	Ala	Gly	Met	Ile	Val	Lys	Phe	Pro	Gly	Thr	Ala	Gln	130	135	140	
Lys	Asn	Thr	Val	Cys	Glu	Pro	Ala	Ser	Pro	Gly	Val	Ser	Pro	Ala	Cys	145	150	155	160
Ala	Ser	Pro	Glu	Asn	Cys	Lys	Glu	Pro	Ser	Ser	Gly	Thr	Ile	Pro	Gln	165	170	175	
Ala	Lys	Pro	Thr	Pro	Val	Ser	Pro	Ala	Thr	Ser	Ser	Ala	Ser	Thr	Met	180	185	190	
Pro	Val	Arg	Gly	Gly	Thr	Arg	Leu	Ala	Gln	Glu	Ala	Ala	Ser	Lys	Leu	195	200	205	
Thr	Arg	Ala	Pro	Asp	Ser	Pro	Ser	Ser	Val	Gly	Arg	Pro	Ser	Ser	Asp	210	215	220	
Pro	Gly	Leu	Ser	Pro	Thr	Gln	Pro	Cys	Pro	Glu	Gly	Ser	Gly	Asp	Cys	225	230	235	240

Arg Lys Gln Cys Glu Pro Asp Tyr Tyr Leu Asp Glu Ala Gly Arg Cys
 245 250 255
 Thr Ala Cys Val Ser Cys Ser Arg Asp Asp Leu Val Glu Lys Thr Pro
 260 265 270
 Cys Ala Trp Asn Ser Ser Arg Thr Cys Glu Cys Arg Pro Gly Met Ile
 275 280 285
 Cys Ala Thr Ser Ala Thr Asn Ser Cys Ala Arg Cys Val Pro Tyr Pro
 290 295 300
 Ile Cys Ala Gly Glu Thr Val Thr Lys Pro Gln Asp Met Ala Glu Lys
 305 310 315 320
 Asp Thr Thr Phe Glu Ala Pro Pro Leu Gly Thr Gln Pro Asp Cys Asn
 325 330 335
 Pro Thr Pro Glu Asn Gly Glu Ala Pro Ala Ser Thr Ser Pro Thr Gln
 340 345 350
 Ser Leu Leu Val Asp Ser Gln Ala Ser Lys Thr Leu Pro Ile Pro Thr
 355 360 365
 Ser Ala Pro Val Ala Leu Ser Ser Thr Gly Lys Pro Val Leu Asp Ala
 370 375 380
 Gly Pro Val Leu Phe Trp Val Ile Leu Val Leu Val Val Val Gly
 385 390 395 400
 Ser Ser Ala Phe Leu Leu Cys His Arg Arg Ala Cys Arg Lys Arg Ile
 405 410 415
 Arg Gln Lys Leu His Leu Cys Tyr Pro Val Gln Thr Ser Gln Pro Lys
 420 425 430
 Leu Glu Leu Val Asp Ser Arg Pro Arg Arg Ser Ser Thr Gln Leu Arg
 435 440 445
 Ser Gly Ala Ser Val Thr Glu Pro Val Ala Glu Glu Arg Gly Leu Met
 450 455 460
 Ser Gln Pro Leu Met Glu Thr Cys His Ser Val Gly Ala Ala Tyr Leu
 465 470 475 480
 Glu Ser Leu Pro Leu Gln Asp Ala Ser Pro Ala Gly Gly Pro Ser Ser
 485 490 495
 Pro Arg Asp Leu Pro Glu Pro Arg Val Ser Thr Glu His Thr Asn Asn
 500 505 510
 Lys Ile Glu Lys Ile Tyr Ile Met Lys Ala Asp Thr Val Ile Val Gly
 515 520 525
 Thr Val Lys Ala Glu Leu Pro Glu Gly Arg Gly Leu Ala Gly Pro Ala
 530 535 540
 Glu Pro Glu Leu Glu Glu Glu Leu Glu Ala Asp His Thr Pro His Tyr
 545 550 555 560

Pro Glu Gln Glu Thr Glu Pro Pro Leu Gly Ser Cys Ser Asp Val Met
565 570 575

Leu Ser Val Glu Glu Glu Gly Lys Glu Asp Pro Leu Pro Thr Ala Ala
580 585 590

Ser Gly Lys
595

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 699 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:
(B) CLONE: hIgG1Fc

(ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 1..696

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAG CCC AGA TCT TGT GAC AAA ACT CAC ACA TGC CCA CCG TGC CCA GCA	48
Glu Pro Arg Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala	
1 5 10 15	
CCT GAA CTC CTG GGG GGA CCG TCA GTC TTC CTC TTC CCC CCA AAA CCC	96
Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro	
20 25 30	
AAG GAC ACC CTC ATG ATC TCC CGG ACC CCT GAG GTC ACA TGC GTG GTG	144
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val	
35 40 45	
GTG GAC GTG AGC CAC GAA GAC CCT GAG GTC AAG TTC AAC TGG TAC GTG	192
Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val	
50 55 60	
GAC GGC GTG GAG GTG CAT AAT GCC AAG ACA AAG CCG CGG GAG GAG CAG	240
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln	
65 70 75 80	
TAC AAC AGC ACG TAC CCG GTG GTC AGC GTC CTC ACC GTC CTG CAC CAG	288
Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln	
85 90 95	

GAC TGG CTG AAT GGC AAG GAC TAC AAG TGC AAG GTC TCC AAC AAA GCC 336
 Asp Trp Leu Asn Gly Lys Asp Tyr Lys Cys Lys Val Ser Asn Lys Ala 110
 100 105 110

CTC CCA GCC CCC ATG CAG AAA ACC ATC TCC AAA GCC AAA GGG CAG CCC 384
 Leu Pro Ala Pro Met Gln Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro 115
 120 125

CGA GAA CCA CAG GTG TAC ACC CTG CCC CCA TCC CGG GAT GAG CTG ACC 432
 Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr 130
 135 140

AAG AAC CAG GTC AGC CTG ACC TGC CTG GTC AAA GGC TTC TAT CCC AGG 480
 Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Arg 145
 150 155 160

CAC ATC GCC GTG GAG TGG GAG AGC AAT GGG CAG CCG GAG AAC AAC TAC 528
 His Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr 165
 170 175

AAG ACC ACG CCT CCC GTG CTG GAC TCC GAC GGC TCC TTC TTC CTC TAC 576
 Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr 180
 185 190

AGC AAG CTC ACC GTG GAC AAG AGC AGG TGG CAG CAG GGG AAC GTC TTC 624
 Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe 195
 200 205

TCA TGC TCC GTG ATG CAT GAG GCT CTG CAC AAC CAC TAC ACG CAG AAG 672
 Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys 210
 215 220

AGC CTC TCC CTG TCT CCG GGT AAA TGA 699
 Ser Leu Ser Leu Ser Pro Gly Lys 225
 230

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 232 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Glu Pro Arg Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
 1 5 10 15

Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
 20 25 30

Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
 35 40 45

Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
 50 55 60

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
 65 70 75 80
 Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
 85 90 95
 Asp Trp Leu Asn Gly Lys Asp Tyr Lys Cys Lys Val Ser Asn Lys Ala
 100 105 110
 Leu Pro Ala Pro Met Gln Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
 115 120 125
 Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr
 130 135 140
 Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Arg
 145 150 155 160
 His Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
 165 170 175
 Lys Thr Thr Pro Val Leu Asp Ser Asp Gly Ser Phe Leu Tyr
 180 185 190
 Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
 195 200 205
 Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
 210 215 220
 Ser Leu Ser Leu Ser Pro Gly Lys
 225 230

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 720 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: muCD30-L
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 1..720

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG GAG CCA GGG CTG CAA CAA GCA GGC AGC TGT GGG GCT CCT TCC CCT 48
 Met Glu Pro Gly Leu Gln Gln Ala Gly Ser Cys Gly Ala Pro Ser Pro
 1 5 10 15

GAC Asp	CCA Pro	GCC Ala	ATG Met	CAG Gln	GTG Val	CAG Gln	CCC Pro	GGC Gly	TCG Ser	GTA Val	GCC Ala	AGC Ser	CCC Pro	TGG Trp	AGA Arg	96
20 25 30																
AGC Ser	ACG Thr	AGG Arg	CCC Pro	TGG Trp	AGA Arg	AGC Ser	ACA Thr	AGT Ser	CGC Arg	AGC Ser	TAC Tyr	TTC Phe	TAC Tyr	CTC Leu	AGC Ser	144
35 40 45																
ACC Thr	ACC Thr	GCA Ala	CTG Leu	GTG Val	TGC Cys	CTT Leu	GTT Val	GTG Val	GCA Ala	GTG Val	GCG Ala	ATC Ile	ATT Ile	CTG Leu	GTA Val	192
50 55 60																
CTG Leu	GTA Val	GTC Val	CAG Gln	AAA Lys	AAG Lys	GAC Asp	TCC Ser	ACT Thr	CCA Pro	AAT Asn	ACA Thr	ACT Thr	GAG Glu	AAG Lys	GCC Ala	240
65 70 75 80																
CCC Pro	CTT Leu	AAA Lys	GGA Gly	GGA Gly	AAT Asn	TGC Cys	TCA Ser	GAG Glu	GAT Asp	CTC Leu	TTC Phe	TGT Cys	ACC Thr	CTG Leu	AAA Lys	288
85 90 95																
AGT Ser	ACT Thr	CCA Pro	TCC Ser	AAG Lys	AAG Lys	TCA Ser	TGG Trp	GCC Tyr	TAC Tyr	CTC Leu	CAA Val	GTG Val	TCA Lys	AAG Lys	CAT His	336
100 105 110																
CTC Leu	AAC Asn	AAT Asn	ACC Thr	AAA Lys	CTG Leu	TCA Ser	TGG Trp	AAC Asn	GAA Glu	GAT Asp	GGC Gly	ACC Ile	ATC His	CAC Gly	GGA Gly	384
115 120 125																
CTC Leu	ATA Ile	TAC Tyr	CAG Gln	GAC Asp	GGG Gly	AAC Asn	CTG Leu	ATA Ile	GTC Val	CAA Gln	TTC Phe	CCT Pro	GGC Gly	TTG Leu	TAC Tyr	432
130 135 140																
TTC Phe	ATC Ile	GTT Val	TGC Cys	CAA Gln	CTG Leu	CAG Gln	TTC Phe	CTC Leu	GTG Val	CAG Val	TGC Cys	TCA Ser	AAT Asn	CAT His	TCT Ser	480
145 150 155 160																
GTG Val	GAC Asp	CTG Leu	ACA Thr	TTG Leu	CAG Gln	CTC Leu	CTC Leu	ATC Ile	AAT Asn	TCC Ser	AAG Lys	ATC Ile	AAA Lys	AAG Lys	CAG Gln	528
165 170 175																
ACG Thr	TTG Leu	GTA Val	ACA Val	GTG Val	TGT Cys	GAG Glu	TCT Ser	GGA Gly	GTT Val	CAG Gln	AGT Ser	AAG Lys	AAC Val	ATC Ile	TAC Tyr	576
180 185 190																
CAG Gln	AAT Asn	CTC Leu	TCT Ser	CAG Gln	TTT Phe	TTG Leu	CTG His	CAT His	TAC Tyr	TTA Leu	CAG Gln	GTC Val	AAC Asn	TCT Ser	ACC Thr	624
195 200 205																
ATA Ile	TCA Ser	GTC Val	AGG Arg	GTG Val	GAT Asp	AAT Asn	TTC Phe	CAG Gln	TAT Tyr	GTG Val	GAT Asp	ACA Thr	AAC Asn	ACT Thr	TTC Phe	672
210 215 220																
CCT Pro	CTT Leu	GAT Asp	AAT Asn	GTG Val	CTA Leu	TCC Ser	GTC Val	TTC Phe	TTA Leu	TAT Tyr	AGT Ser	AGC Ser	TCA Ser	GAC Asp	TGA Asp	720
225 230 235 240																

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 239 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Glu Pro Gly Leu Gln Gln Ala Gly Ser Cys Gly Ala Pro Ser Pro
1 5 10 15
Asp Pro Ala Met Gln Val Gln Pro Gly Ser Val Ala Ser Pro Trp Arg
20 25 30
Ser Thr Arg Pro Trp Arg Ser Thr Ser Arg Ser Tyr Phe Tyr Leu Ser
35 40 45
Thr Thr Ala Leu Val Cys Leu Val Val Ala Val Ala Ile Ile Leu Val
50 55 60
Leu Val Val Gln Lys Lys Asp Ser Thr Pro Asn Thr Thr Glu Lys Ala
65 70 75 80
Pro Leu Lys Gly Gly Asn Cys Ser Glu Asp Leu Phe Cys Thr Leu Lys
85 90 95
Ser Thr Pro Ser Lys Lys Ser Trp Ala Tyr Leu Gln Val Ser Lys His
100 105 110
Leu Asn Asn Thr Lys Leu Ser Trp Asn Glu Asp Gly Thr Ile His Gly
115 120 125
Leu Ile Tyr Gln Asp Gly Asn Leu Ile Val Gln Phe Pro Gly Leu Tyr
130 135 140
Phe Ile Val Cys Gln Leu Gln Phe Leu Val Gln Cys Ser Asn His Ser
145 150 155 160
Val Asp Leu Thr Leu Gln Leu Leu Ile Asn Ser Lys Ile Lys Lys Gln
165 170 175
Thr Leu Val Thr Val Cys Glu Ser Gly Val Gln Ser Lys Asn Ile Tyr
180 185 190
Gln Asn Leu Ser Gln Phe Leu Leu His Tyr Leu Gln Val Asn Ser Thr
195 200 205
Ile Ser Val Arg Val Asp Asn Phe Gln Tyr Val Asp Thr Asn Thr Phe
210 215 220
Pro Leu Asp Asn Val Leu Ser Val Phe Leu Tyr Ser Ser Ser Asp
225 230 235

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 705 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(vii) IMMEDIATE SOURCE:
 (B) CLONE: huCD30-L

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..705

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATG GAC CCA GGG CTG CAG CAA GCA CTC AAC GGA ATG GCC CCT CCT GGA	48
Met Asp Pro Gly Leu Gln Gln Ala Leu Asn Gly Met Ala Pro Pro Gly	
1 5 10 15	
GAC ACA GCC ATG CAT GTG CCG GCG GGC TCC GTG GCC AGC CAC CTG GGG	96
Asp Thr Ala Met His Val Pro Ala Gly Ser Val Ala Ser His Leu Gly	
20 25 30	
ACC ACG AGC CGC AGC TAT TTC TAT TTG ACC ACA GCC ACT CTG GCT CTG	144
Thr Thr Ser Arg Ser Tyr Phe Tyr Leu Thr Thr Ala Thr Leu Ala Leu	
35 40 45	
TGC CTT GTC TTC ACG GTG GCC ACT ATT ATG GTG TTG GTC GTT CAG AGG	192
Cys Leu Val Phe Thr Val Ala Thr Ile Met Val Leu Val Val Gln Arg	
50 55 60	
ACG GAC TCC ATT CCC AAC TCA CCT GAC AAC GTC CCC CTC AAA GGA GGA	240
Thr Asp Ser Ile Pro Asn Ser Pro Asp Asn Val Pro Leu Lys Gly Gly	
65 70 75 80	
AAT TGC TCA GAA GAC CTC TTA TGT ATC CTG AAA AGA GCT CCA TTC AAG	288
Asn Cys Ser Glu Asp Leu Leu Cys Ile Leu Lys Arg Ala Pro Phe Lys	
85 90 95	
AAG TCA TGG GCC TAC CTC CAA GTG GCA AAG CAT CTA AAC AAA ACC AAG	336
Lys Ser Trp Ala Tyr Leu Gln Val Ala Lys His Leu Asn Lys Thr Lys	
100 105 110	
TTG TCT TGG AAC AAA GAT GGC ATT CTC CAT GGA GTC AGA TAT CAG GAT	384
Leu Ser Trp Asn Lys Asp Gly Ile Leu His Gly Val Arg Tyr Gln Asp	
115 120 125	
GGG AAT CTG GTG ATC CAA TTC CCT GGT TTG TAC TTC ATC ATT TGC CAA	432
Gly Asn Leu Val Ile Gln Phe Pro Gly Leu Tyr Phe Ile Ile Cys Gln	
130 135 140	
CTG CAG TTT CTT GTA CAA TGC CCA AAT AAT TCT GTC GAT CTG AAG TTG	480
Leu Gln Phe Leu Val Gln Cys Pro Asn Asn Ser Val Asp Leu Lys Leu	
145 150 155 160	

GAG CTT CTC ATC AAC AAG CAT ATC AAA AAA CAG GCC CTG GTG ACA GTG 528
 Glu Leu Leu Ile Asn Lys His Ile Lys Lys Gln Ala Leu Val Thr Val 175
 165 170 175

TGT GAG TCT GGA ATG CAA ACG AAA CAC GTA TAC CAG AAT CTC TCT CAA 576
 Cys Glu Ser Gly Met Gln Thr Lys His Val Tyr Gln Asn Leu Ser Gln 190
 180 185 190

TTC TTG CTG GAT TAC CTG CAG GTC AAC ACC ACC ATA TCA GTC AAT GTG 624
 Phe Leu Leu Asp Tyr Leu Gln Val Asn Thr Thr Ile Ser Val Asn Val 205
 195 200 205

GAT ACA TTC CAG TAC ATA GAT ACA AGC ACC TTT CCT CTT GAG AAT GTG 672
 Asp Thr Phe Gln Tyr Ile Asp Thr Ser Thr Phe Pro Leu Glu Asn Val 220
 210 215 220

TTG TCC ATC TTC TTA TAC AGT AAT TCA GAC TGA 705
 Leu Ser Ile Phe Leu Tyr Ser Asn Ser Asp 235
 225 230 235

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 234 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Asp Pro Gly Leu Gln Gln Ala Leu Asn Gly Met Ala Pro Pro Gly
 1 5 10 15

Asp Thr Ala Met His Val Pro Ala Gly Ser Val Ala Ser His Leu Gly
 20 25 30

Thr Thr Ser Arg Ser Tyr Phe Tyr Leu Thr Thr Ala Thr Leu Ala Leu
 35 40 45

Cys Leu Val Phe Thr Val Ala Thr Ile Met Val Leu Val Val Gln Arg
 50 55 60

Thr Asp Ser Ile Pro Asn Ser Pro Asp Asn Val Pro Leu Lys Gly Gly
 65 70 75 80

Asn Cys Ser Glu Asp Leu Leu Cys Ile Leu Lys Arg Ala Pro Phe Lys
 85 90 95

Lys Ser Trp Ala Tyr Leu Gln Val Ala Lys His Leu Asn Lys Thr Lys
 100 105 110

Leu Ser Trp Asn Lys Asp Gly Ile Leu His Gly Val Arg Tyr Gln Asp
 115 120 125

Gly Asn Leu Val Ile Gln Phe Pro Gly Leu Tyr Phe Ile Ile Cys Gln
 130 135 140

Leu Gln Phe Leu Val Gln Cys Pro Asn Asn Ser Val Asp Leu Lys Leu
 145 150 155 160
 Glu Leu Leu Ile Asn Lys His Ile Lys Lys Gln Ala Leu Val Thr Val
 165 170 175
 Cys Glu Ser Gly Met Gln Thr Lys His Val Tyr Gln Asn Leu Ser Gln
 180 185 190
 Phe Leu Leu Asp Tyr Leu Gln Val Asn Thr Thr Ile Ser Val Asn Val
 195 200 205
 Asp Thr Phe Gln Tyr Ile Asp Thr Ser Thr Phe Pro Leu Glu Asn Val
 210 215 220
 Leu Ser Ile Phe Leu Tyr Ser Asn Ser Asp
 225 230

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: 5' PCR Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATAGCGGCCG CCACCATGCG CGTCCTCCTC GCCGCGCTG

39

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: 3' PCR Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ACAAGATCTG GGCTCCTTCC CCGTGGAGGA GAGAGCGAC

39

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(iii) MOLECULE TYPE: DNA

(vii) IMMEDIATE SOURCE:
(B) CLONE: BGL II Adaptor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GATCTGGCAA CGAAGGTACC ATGG

24

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(vii) IMMEDIATE SOURCE:
(B) CLONE: BGL II Adaptor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CCATGGTACC TTCGTTGCCA

20

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 33 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vii) IMMEDIATE SOURCE:
(B) CLONE: Upstream sequence

(ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 1..33

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

ATG GGC TGT GGG GCT CCT TCC CCT GAC CCA GCC
Met Gly Cys Gly Ala Pro Ser Pro Asp Pro Ala
1 5 10

33

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met Gly Cys Gly Ala Pro Ser Pro Asp Pro Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:
(B) CLONE: FLAG peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Asp Tyr Lys Asp Asp Asp Asp Lys
1 5

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(vii) IMMEDIATE SOURCE:
(B) CLONE: Murine cDNA Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

AGATGCTTTG ACACTTG

17

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (vii) IMMEDIATE SOURCE:
(B) CLONE: Human cDNA Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

ATCACCAGAT TCCCATC

17

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 663 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (vii) IMMEDIATE SOURCE:
(B) CLONE: muCD30-L

- (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 1..663

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

ATG	CAG	GTG	CAG	CCC	GGC	TCG	GTA	GCC	AGC	CCC	TGG	AGA	AGC	ACG	AGG	48
Met	Gln	Val	Gln	Pro	Gly	Ser	Val	Ala	Ser	Pro	Trp	Arg	Ser	Thr	Arg	
1				5					10					15		

CCC	TGG	AGA	AGC	ACA	AGT	CGC	AGC	TAC	TTC	TAC	CTC	AGC	ACC	ACC	GCA	96
Pro	Trp	Arg	Ser	Thr	Ser	Arg	Ser	Tyr	Phe	Tyr	Leu	Ser	Thr	Thr	Ala	
			20					25					30			

CTG	GTG	TGC	CTT	GTT	GTG	GCA	GTG	GCG	ATC	ATT	CTG	GTA	CTG	GTA	GTC	144
Leu	Val	Cys	Leu	Val	Val	Ala	Val	Ala	Ile	Ile	Leu	Val	Leu	Val	Val	
		35					40					45				

CAG AAA AAG GAC TCC ACT CCA AAT ACA ACT GAG AAG GCC CCC CTT AAA	192
Gln Lys Lys Asp Ser Thr Pro Asn Thr Thr Glu Lys Ala Pro Leu Lys	
50 55 60	
GGA GGA AAT TGC TCA GAG GAT CTC TTC TGT ACC CTG AAA AGT ACT CCA	240
Gly Gly Asn Cys Ser Glu Asp Leu Phe Cys Thr Leu Lys Ser Thr Pro	
65 70 75 80	
TCC AAG AAG TCA TGG GCC TAC CTC CAA GTG TCA AAG CAT CTC AAC AAT	288
Ser Lys Lys Ser Trp Ala Tyr Leu Gln Val Ser Lys His Leu Asn Asn	
85 90 95	
ACC AAA CTG TCA TGG AAC GAA GAT GGC ACC ATC CAC GGA CTC ATA TAC	336
Thr Lys Leu Ser Trp Asn Glu Asp Gly Thr Ile His Gly Leu Ile Tyr	
100 105 110	
CAG GAC GGG AAC CTG ATA GTC CAA TTC CCT GGC TTG TAC TTC ATC GTT	384
Gln Asp Gly Asn Leu Ile Val Gln Phe Pro Gly Leu Tyr Phe Ile Val	
115 120 125	
TGC CAA CTG CAG TTC CTC GTG CAG TGC TCA AAT CAT TCT GTG GAC CTG	432
Cys Gln Leu Gln Phe Leu Val Gln Cys Ser Asn His Ser Val Asp Leu	
130 135 140	
ACA TTG CAG CTC CTC ATC AAT TCC AAG ATC AAA AAG CAG ACG TTG GTA	480
Thr Leu Gln Leu Leu Ile Asn Ser Lys Ile Lys Lys Gln Thr Leu Val	
145 150 155 160	
ACA GTG TGT GAG TCT GGA GTT CAG AGT AAG AAC ATC TAC CAG AAT CTC	528
Thr Val Cys Glu Ser Gly Val Gln Ser Lys Asn Ile Tyr Gln Asn Leu	
165 170 175	
TCT CAG TTT TTG CTG CAT TAC TTA CAG GTC AAC TCT ACC ATA TCA GTC	576
Ser Gln Phe Leu Leu His Tyr Leu Gln Val Asn Ser Thr Ile Ser Val	
180 185 190	
AGG GTG GAT AAT TTC CAG TAT GTG GAT ACA AAC ACT TTC CCT CTT GAT	624
Arg Val Asp Asn Phe Gln Tyr Val Asp Thr Asn Thr Phe Pro Leu Asp	
195 200 205	
AAT GTG CTA TCC GTC TTC TTA TAT AGT AGC TCA GAC TGA	663
Asn Val Leu Ser Val Phe Leu Tyr Ser Ser Ser Asp	
210 215 220	

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 220 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Met Gln Val Gln Pro Gly Ser Val Ala Ser Pro Trp Arg Ser Thr Arg
1 5 10 15

Pro Trp Arg Ser Thr Ser Arg Ser Tyr Phe Tyr Leu Ser Thr Thr Ala
 20 25 30
 Leu Val Cys Leu Val Val Ala Val Ala Ile Ile Leu Val Leu Val Val
 35 40 45
 Gln Lys Lys Asp Ser Thr Pro Asn Thr Thr Glu Lys Ala Pro Leu Lys
 50 55 60
 Gly Gly Asn Cys Ser Glu Asp Leu Phe Cys Thr Leu Lys Ser Thr Pro
 65 70 75 80
 Ser Lys Lys Ser Trp Ala Tyr Leu Gln Val Ser Lys His Leu Asn Asn
 85 90 95
 Thr Lys Leu Ser Trp Asn Glu Asp Gly Thr Ile His Gly Leu Ile Tyr
 100 105 110
 Gln Asp Gly Asn Leu Ile Val Gln Phe Pro Gly Leu Tyr Phe Ile Val
 115 120 125
 Cys Gln Leu Gln Phe Leu Val Gln Cys Ser Asn His Ser Val Asp Leu
 130 135 140
 Thr Leu Gln Leu Leu Ile Asn Ser Lys Ile Lys Lys Gln Thr Leu Val
 145 150 155 160
 Thr Val Cys Glu Ser Gly Val Gln Ser Lys Asn Ile Tyr Gln Asn Leu
 165 170 175
 Ser Gln Phe Leu Leu His Tyr Leu Gln Val Asn Ser Thr Ile Ser Val
 180 185 190
 Arg Val Asp Asn Phe Gln Tyr Val Asp Thr Asn Thr Phe Pro Leu Asp
 195 200 205
 Asn Val Leu Ser Val Phe Leu Tyr Ser Ser Ser Ser Asp
 210 215 220

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 125 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein fragment
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vii) IMMEDIATE SOURCE:
 - (B) CLONE: huCD30 fragment (PRELIM)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Pro Gly Asp Thr Val Xaa His Val Pro Ala Gly Ser Glu Ala Ser His
 1 5 10 15
 Leu Gly Thr Thr Ser Arg Xaa Tyr Phe Tyr Leu Thr Thr Xaa Thr Leu
 20 25 30
 Ala Leu Cys Leu Val Phe Thr Val Ala Thr Ile Met Val Leu Val Val
 35 40 45
 Gln Arg Thr Asp Ser Ile Pro Asn Ser Pro Asp Asn Val Pro Leu Lys
 50 55 60
 Gly Gly Asn Cys Ser Glu Asp Leu Leu Cys Ile Leu Lys Arg Ala Pro
 65 70 75 80
 Phe Lys Lys Ser Trp Ala Tyr Leu Gln Val Xaa Lys His Leu Asn Lys
 85 90 95
 Thr Xaa Leu Ser Trp Asn Lys Asp Gly Ile Leu His Gly Val Arg Tyr
 100 105 110
 Gln Asp Gly Asn Leu Val Ile Gln Phe Pro Gly Phe Val
 115 120 125

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 130 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein fragment

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

- (B) CLONE: muCD30 fragment (PRELIM)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Met Gln Val Gln Pro Gly Ser Val Ala Ser Pro Trp Arg Ser Thr Arg
 1 5 10 15
 Pro Trp Arg Ser Thr Ser Arg Ser Tyr Phe Tyr Leu Ser Thr Thr Ala
 20 25 30
 Leu Val Cys Leu Val Val Xaa Val Ala Ile Ile Leu Val Leu Val Val
 35 40 45
 Gln Lys Lys Asp Ser Thr Pro Asn Thr Thr Glu Lys Ala Pro Leu Lys
 50 55 60
 Gly Gly Asn Cys Ser Glu Asp Leu Phe Cys Thr Leu Lys Ser Thr Pro
 65 70 75 80

Ser Lys Lys Ser Trp Ala Tyr Leu Gln Val Ser Lys His Leu Asn Asn 85 90 95
 Thr Lys Leu Ser Trp Asn Glu Asp Gly Thr Ile His Gly Leu Ile Tyr 100 105 110
 Gln Asp Gly Asn Leu Ile Val Gln Phe Pro Gly Leu Tyr Phe Ile Val 115 120 125
 Cys Gln 130

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 648 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(vii) IMMEDIATE SOURCE:
 (B) CLONE: huCD30-L

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..648

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

ATG CAT GTG CCG GCG GGC TCC GTG GCC AGC CAC CTG GGG ACC ACG AGC	48
Met His Val Pro Ala Gly Ser Val Ala Ser His Leu Gly Thr Thr Ser	
1 5 10 15	
CGC AGC TAT TTC TAT TTG ACC ACA GCC ACT CTG GCT CTG TGC CTT GTC	96
Arg Ser Tyr Phe Tyr Leu Thr Thr Ala Thr Leu Ala Leu Cys Leu Val	
20 25 30	
TTC ACG GTG GCC ACT ATT ATG GTG TTG GTC GTT CAG AGG ACG GAC TCC	144
Phe Thr Val Ala Thr Ile Met Val Leu Val Val Gln Arg Thr Asp Ser	
35 40 45	
ATT CCC AAC TCA CCT GAC AAC GTC CCC CTC AAA GGA GGA AAT TGC TCA	192
Ile Pro Asn Ser Pro Asp Asn Val Pro Leu Lys Gly Gly Asn Cys Ser	
50 55 60	
GAA GAC CTC TTA TGT ATC CTG AAA AGA GCT CCA TTC AAG AAG TCA TGG	240
Glu Asp Leu Leu Cys Ile Leu Lys Arg Ala Pro Phe Lys Lys Ser Trp	
65 70 75 80	
GCC TAC CTC CAA GTG GCA AAG CAT CTA AAC AAA ACC AAG TTG TCT TGG	288
Ala Tyr Leu Gln Val Ala Lys His Leu Asn Lys Thr Lys Leu Ser Trp	
85 90 95	

AAC AAA GAT GGC ATT CTC CAT GGA GTC AGA TAT CAG GAT GGG AAT CTG 336
Asn Lys Asp Gly Ile Leu His Gly Val Arg Tyr Gln Asp Gly Asn Leu
100 105 110
GTG ATC CAA TTC CCT GGT TTG TAC TTC ATC ATT TGC CAA CTG CAG TTT 384
Val Ile Gln Phe Pro Gly Leu Tyr Phe Ile Ile Cys Gln Leu Gln Phe
115 120 125
CTT GTA CAA TGC CCA AAT AAT TCT GTC GAT CTG AAG TTG GAG CTT CTC 432
Leu Val Gln Cys Pro Asn Asn Ser Val Asp Leu Lys Leu Glu Leu Leu
130 135 140
ATC AAC AAG CAT ATC AAA AAA CAG GCC CTG GTG ACA GTG TGT GAG TCT 480
Ile Asn Lys His Ile Lys Lys Gln Ala Leu Val Thr Val Cys Glu Ser
145 150 155 160
GGA ATG CAA ACG AAA CAC GTA TAC CAG AAT CTC TCT CAA TTC TTG CTG 528
Gly Met Gln Thr Lys His Val Tyr Gln Asn Leu Ser Gln Phe Leu Leu
165 170 175
GAT TAC CTG CAG GTC AAC ACC ACC ATA TCA GTC AAT GTG GAT ACA TTC 576
Asp Tyr Leu Gln Val Asn Thr Thr Ile Ser Val Asn Val Asp Thr Phe
180 185 190
CAG TAC ATA GAT ACA AGC ACC TTT CCT CTT GAG AAT GTG TTG TCC ATC 624
Gln Tyr Ile Asp Thr Ser Thr Phe Pro Leu Glu Asn Val Leu Ser Ile
195 200 205
TTC TTA TAC AGT AAT TCA GAC TGA 648
Phe Leu Tyr Ser Asn Ser Asp
210 215

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 215 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Met His Val Pro Ala Gly Ser Val Ala Ser His Leu Gly Thr Thr Ser
1 5 10 15
Arg Ser Tyr Phe Tyr Leu Thr Thr Ala Thr Leu Ala Leu Cys Leu Val
20 25 30
Phe Thr Val Ala Thr Ile Met Val Leu Val Val Gln Arg Thr Asp Ser
35 40 45
Ile Pro Asn Ser Pro Asp Asn Val Pro Leu Lys Gly Asn Cys Ser
50 55 60
Glu Asp Leu Leu Cys Ile Leu Lys Arg Ala Pro Phe Lys Lys Ser Trp
65 70 75 80

Ala Tyr Leu Gln Val Ala Lys His Leu Asn Lys Thr Lys Leu Ser Trp
 85 90 95
 Asn Lys Asp Gly Ile Leu His Gly Val Arg Tyr Gln Asp Gly Asn Leu
 100 105 110
 Val Ile Gln Phe Pro Gly Leu Tyr Phe Ile Ile Cys Gln Leu Gln Phe
 115 120 125
 Leu Val Gln Cys Pro Asn Asn Ser Val Asp Leu Lys Leu Glu Leu Leu
 130 135 140
 Ile Asn Lys His Ile Lys Lys Gln Ala Leu Val Thr Val Cys Glu Ser
 145 150 155 160
 Gly Met Gln Thr Lys His Val Tyr Gln Asn Leu Ser Gln Phe Leu Leu
 165 170 175
 Asp Tyr Leu Gln Val Asn Thr Thr Ile Ser Val Asn Val Asp Thr Phe
 180 185 190
 Gln Tyr Ile Asp Thr Ser Thr Phe Pro Leu Glu Asn Val Leu Ser Ile
 195 200 205
 Phe Leu Tyr Ser Asn Ser Asp
 210 215